



## **Effect of Integrated Nutrient Management on Plant Growth and Tuber Yield of Dahlia (*Dahlia variabilis*) cv. Kenya White**

**T. Raghunandan<sup>a\*</sup>, J. P. Collis<sup>a</sup>, S. Saravanan<sup>a</sup> and K. S. Barman<sup>a</sup>**

<sup>a</sup> Allahabad School of Agriculture, SHIATS, Allahabad, Uttar Pradesh, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJPSS/2021/v33i2430753

#### Editor(s):

- (1) Dr. Hakan Sevik, Kastamonu University, Turkey.  
(2) Dr. Hon H. Ho, State University of New York, USA.

#### Reviewers:

- (1) Gyula Oros, Plant Protection Institute HAS, India.  
(2) Mohssen Hassan Mohamed Elbagory, King Khalid University, KSA & Soil, Water and Environment Research Institute (SWERI), Agricultural Research Center (ARC), Egypt.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:  
<https://www.sdiarticle5.com/review-history/70294>

**Original Research Article**

**Received 28 June 2021**  
**Accepted 02 August 2021**  
**Published 13 December 2021**

### **ABSTRACT**

The experiment was carried out at Experimental field, Department of Horticulture, Sam Higginbottom Institute of Agriculture Technology and Sciences [formerly known as Allahabad Agriculture Institute Deemed University, AAI-DU] during the Spring season 2014-2015. The experiment consists of Thirteen treatments viz., (T<sub>0</sub>) Control-RDF (100:120:100 N P K kg ha<sup>-1</sup>), (T<sub>1</sub>) 75% RDF + AZ @ 2.5 kg ha<sup>-1</sup>, (T<sub>2</sub>) 75% RDF + FYM @ 10 ton ha<sup>-1</sup>, (T<sub>3</sub>) 75% RDF + VC @ 10 ton ha<sup>-1</sup>, (T<sub>4</sub>) 75% RDF + AZ @ 2.5 kg/ha +FYM @ 10 ton ha<sup>-1</sup>+VC @ 10 ton ha<sup>-1</sup>, (T<sub>5</sub>) 50% RDF +AZ @ 2.5 kg ha<sup>-1</sup>, (T<sub>6</sub>) 50% RDF + FYM@ 10 ton ha<sup>-1</sup>, (T<sub>7</sub>) 50% RDF + VC@ 10 ton ha<sup>-1</sup>, (T<sub>8</sub>) 50% RDF + AZ @ 2.5 kg ha<sup>-1</sup>+FYM @ 10 ton ha<sup>-1</sup>+VC @ 10 ton ha<sup>-1</sup>, (T<sub>9</sub>) 25% RDF + AZ@ 2.5 kg ha<sup>-1</sup>, (T<sub>10</sub>) 25% RDF + FYM@ 10 ton ha<sup>-1</sup>, (T<sub>11</sub>) 25% RDF + VC@ 10 ton ha<sup>-1</sup>, (T<sub>12</sub>) 25% RDF + AZ @ 2.5 kg ha<sup>-1</sup>+FYM @ 10 ton ha<sup>-1</sup>+VC @ 10 ton ha<sup>-1</sup>. The treatments were replicated thrice in a Randomized Complete Block Design. The results revealed that Plants treated with 75% RDF + AZ @ 2.5 kg/ha +FYM @ 10 t/ha +VC @ 10 t/ha (T<sub>4</sub>) significantly recorded maximum vegetative and Tuber yield attributes like plant height (91.87 cm), plant spread (92.38 cm), Number of Branches per plant (7.27), number of leaves per plant (26.53), number of tubers per plant (4.80), maximum tuber weight (958.53) and Maximum tuber yield per plant was recorded in T<sub>4</sub> (75% RDF + AZ @ 2.5 kg/ha +FYM @ 10 t/ha +VC @ 10 t/ha) (43.33g) followed by T<sub>3</sub> (75% RDF + VC @ 10 t/ha) (40.95g).

\*Corresponding author: E-mail: [raghunandanthogari@gmail.com](mailto:raghunandanthogari@gmail.com);

**Keywords:** *Dahlia; Kenya white; tuber yield; RDF (Recommended Dose of Fertilizer); azotobacter; vermicompost; FYM.*

## 1. INTRODUCTION

Dahlia is one of the most popular bulbous flower crop grown in the world for its beautiful ornamental blooms of various shades of colours for the beautification of gardens and cut flowers. It is a tuberous rooted, half-hardy herbaceous perennial belonging to the family Asteraceae, having its origin in Mexico. Dahlia occupies a place of pride in any garden and they are grown both in field and pot culture and are extensively used for exhibition, garden display and home decoration. For exhibition and garden display all types of dahlias are used. Dwarf growing types are suitable for beds and borders (pure/mixed borders). Large flowering dahlias in pots are popular for terrace garden or verandah display [1]. There are so many factors responsible for growth and development of plants such as soil type, prevailing climatic conditions of an area, nutritional factors, irrigation, plant density per unit area, season of growing, etc. among them Nutrition plays a major role in growth, flowering and yield of flower crop. Integrated nutrient management refers to the maintenance of soil fertility and of plant nutrient supply at an optimum level for sustaining the desired productivity through optimization of the benefits from all possible sources of organic, inorganic and biological components in an integrated manner [2]. Application of nutrients from organic as well as inorganic sources. It aims at improving their physico-chemical and biological properties of the soil. Integrated Nutrient Management refers to regulated nutrient supply for optimum crop growth and higher productivity, improvement and maintenance of soil fertility, Zero adverse impact on agro-ecosystem quality by balanced fertilization of organic manures, inorganic fertilizers and bio-inoculants. Keeping the above facts in view, an experiment was conducted "Effect of integrated nutrient management on plant growth and tuber yield of Dahlia (*Dahlia variabilis*) cv. Kenya white under Allahabad agro climatic conditions." Prasad et al. [3].

## 2. MATERIALS AND METHODS

The present experiment was carried out at Experimental field, Department of Horticulture, Sam Higginbottom Institute of Agriculture Technology and Sciences [formerly known as Allahabad Agriculture Institute Deemed University, AAI-DU] during the Spring

season 2014-2015. The experiment consists of Thirteen treatments viz., (T0) Control-RDF (100:120:100 N P K kg ha<sup>-1</sup>), (T1) 75% RDF + AZ@ 2.5 kg ha<sup>-1</sup>, (T2) 75% RDF + FYM @ 10 ton ha<sup>-1</sup>, (T3) 75% RDF + VC@ 10 ton ha<sup>-1</sup>, (T4) 75% RDF + AZ @ 2.5 kg/ha +FYM @ 10 ton ha<sup>-1</sup>+VC @ 10 ton ha<sup>-1</sup>, (T5) 50% RDF +AZ@ 2.5 kg ha<sup>-1</sup>, (T6) 50% RDF + FYM@ 10 ton ha<sup>-1</sup>, (T7) 50% RDF + VC@ 10 ton ha<sup>-1</sup>, (T8) 50% RDF + AZ @ 2.5 kg ha<sup>-1</sup>+FYM @ 10 ton ha<sup>-1</sup>+VC @ 10 ton ha<sup>-1</sup>, (T9) 25% RDF + AZ@ 2.5 kg ha<sup>-1</sup>, (T10) 25% RDF + FYM@ 10 ton ha<sup>-1</sup>, (T11) 25% RDF + VC@ 10 ton ha<sup>-1</sup>+, (T12) 25% RDF + AZ @ 2.5 kg ha<sup>-1</sup>+FYM @ 10 ton ha<sup>-1</sup>+VC @ 10 ton ha<sup>-1</sup>. The treatments were replicated thrice in a Randomized Complete Block Design Prasad et al. [3]. Organic Manure and fertilizers (NPK) were applied according to recommended doses for dahlia, i.e. Urea: SSP: MOP @ 100: 120: 100 kg ha<sup>-1</sup>. Organic manure viz., FYM and Vermicompost was well incorporated in the experimental field before transplanting the seedlings in advance. 40 days old healthy and uniformly grown seedlings were transplanted with a spacing of 45cm x 60 cm. Nitrogen, phosphorous and potassium were applied in the form of Urea, Single Super Phosphate (SSP) and Muriate of potash (MOP), respectively. At the time of transplanting, half of the dose of N and full doses of phosphorous and potassium was applied and remaining two doses of Nitrogen was applied after 30 days transplanting. All the growth and tuber yield parameters were recorded timely and the data were analyzed in Randomized Complete Block Design (RCBD) as per Gomez and Gomez [1].

## 3. RESULTS AND DISCUSSION

### 3.1 Growth Parameters

From the data presented in Table 1 clearly shows that all the growth parameters were significantly influenced by different treatment combinations. The growth parameters like plant height (91.87 cm), plant spread (92.38 cm), number of Branches per plant (7.27) and number of leaves per plant (26.53) were recorded maximum in T4 (75%RDF+ AZ @ 2.5 kg/ha +FYM @ 10 t/ha +VC @ 10 t/ha) whereas, all the growth parameters were recorded minimum in T0, i.e. control Wararkar et al [4]. The results clearly shows that combined application of nutrients viz., 75% RDF, Azotobacter, Farm Yard Manure (FYM) and Vermicompost (VC)

**Table 1. Effect of Integrated Nutrient Management on Growth and Tuber yield parameters of Dahlia (*Dahlia variabilis*) cv. Kenya white**

Treatments	Plant Height (cm)	Plant Spread (cm)	Number of Branches Per plant	Number of leaves	Tuber weight (g)	Number of tubers per plant	Tuber Yield Per plant
T <sub>0</sub> Control-RDF (100:120:100 N P K Kg/ha)	67.03	81.07	6.47	19.20	193.33	2.47	29.52
T <sub>1</sub> 75% RDF + AZ@ 2.5 kg/ha	85.93	89.64	6.87	23.20	849.00	3.87	38.28
T <sub>2</sub> 75% RDF + FYM @ 10 t/ha	86.43	90.76	7.00	23.80	899.00	4.00	38.57
T <sub>3</sub> 75% RDF + VC@ 10 t/ha	89.90	91.87	7.00	25.27	941.67	4.40	40.95
T <sub>4</sub> 75% RDF + AZ @ 2.5 kg/ha +FYM @ 10 t/ha +VC @ 10t/ha	91.87	92.38	7.27	26.53	958.53	4.80	43.33
T <sub>5</sub> 50% RDF +AZ@ 2.5 kg/ha	75.30	85.57	6.67	21.00	461.33	3.93	33.80
T <sub>6</sub> 50% RDF + FYM@ 10 t/ha	82.00	88.01	6.60	22.87	650.00	3.67	35.71
T <sub>7</sub> 50% RDF + VC@ 10 t/ha	84.27	89.61	6.47	22.80	716.00	3.93	38.28
T <sub>8</sub> 50% RDF + AZ @ 2.5 kg/ha +FYM @ 10 t/ha +VC @ 10 t/ha	88.83	90.13	6.87	24.80	923.87	4.53	39.24
T <sub>9</sub> 25% RDF + AZ@ 2.5 kg/ha	70.70	83.23	6.67	20.53	233.00	3.40	30.00
T <sub>10</sub> 25% RDF + FYM@ 10 t/ha	71.80	85.06	6.67	20.40	346.67	3.53	33.04
T <sub>11</sub> 25% RDF + VC@ 10 t/ha	76.27	86.46	6.93	20.80	524.40	3.53	34.28
T <sub>12</sub> 25% RDF + AZ @ 2.5 kg/ha +FYM @ 10 t/ha +VC @ 10 t/ha	79.67	86.87	7.07	21.87	558.53	4.00	35.71
F- test	S	S	S	S	S	S	S
S. Ed. (±)	0.340	0.187	0.148	0.303	2.537	0.136	1.492
C. D. (P = 0.05)	0.702	0.386	0.305	0.624	5.236	0.281	3.080

enhanced the vigorous growth of the plant. It might be due to presence of more readily available forms of nutrients i.e., nitrogen, phosphorous, potassium in combination with organic sources viz., Azotobacter, FYM and VC helped in fixing atmospheric nitrogen and solubilized phosphorous. Application of Azotobacter with RDF might also secretes some growth promoters viz., auxin, gibberellins, organic acids and vitamins, in soil with bio inoculation which increases the photosynthetic activity of the plant and better flow of essential nutrients leading to better growth of the plant these results are in conformity with Kulkarni [2] in aster, Chandrikapure et al, [5] in marigold and Kumar et al. [6] in aster. Prasad et al. [3].

### 3.2 Tuberos Yield Parameters

The data revealed that different treatment combination affected all the tuberos yield parameters as shown in Table 1. The maximum tuber weight (958.53g), number of tubers per plant (4.80) and maximum tuber yield per plant was recorded in T<sub>4</sub> (75% RDF + AZ @ 2.5 kg/ha +FYM @ 10 t/ha +VC @ 10 t/ha) (43.33g) followed by T<sub>3</sub> (75% RDF + VC @ 10 t/ha) (40.95g). The increase in tuber weight, number of tubers/plant Prasad et al. [3], tuber yield might be due to recommended dose of fertilizers application along with Azotobacter and Vermicompost, where cell enlargement and increased production of carbohydrate synthesis may be caused by Azotobacter which transferred into tuber development. Recommended dose of Fertilizers application made readily available form of nutrients to plants, in which phosphorous helped in easy proliferation and better root formation which lead to increase in tuber yield parameters. These results are in conformity with Kumar et al [7], Wararkar et al [4], Pandey et al, [8], Sabah et al. [9] and Sheergojri et al [10].

### 4. CONCLUSION

From the above investigation, it can be concluded that integrated approach i.e., application of Recommended dose of fertilizers along with organic manures and biofertilizers showed significant improvement in all growth and yield parameters of Dahlia var. Kenya White. Among the different treatment combination applied, Treatment T<sub>4</sub> (75% RDF + AZ @ 2.5 kg/ha +FYM @ 10 t/ha +VC @ 10 t/ha) found best with respect to growth parameters where as maximum tuber weight, number of tubers per plant and maximum tuber yield per plant was

recorded in T<sub>4</sub> (75% RDF + AZ @ 2.5 kg/ha +FYM @ 10 t/ha +VC @ 10 t/ha) followed by T<sub>3</sub> (75% RDF + VC @ 10 t/ha) (40.95g).

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research, John Wiley and Sons. 1984;680.
2. Kulkarni RG. Studies on effect of Azotobacter and Azospirillum alone and in combination under graded levels of nitrogen on growth and yield Aster. M.Sc. Thesis submitted to M.P.K.V., Rahuri; 1990.
3. Prasad DS, Prasad VM, Goutham SK, Bose SC. Effect of integrated nutrient management on flowering and flower yield of dahlia (*Dahlia variabilis* L) cv. Kenya Orange. Plant Archives. 2018;18(1):795-8.
4. Wararkar SM, Sarvanan S, Prasad VM. Effect of integrated nutrient management on growth, flowering and yield of dahlia (*dahlia variabilis* l.). Cv. Kenya white. Plant Archives. 2020;20(1):3292-6.
5. Chandrikapure KR, Sadavarte KT, Panchabhai DM, Shelke BD. Effect of bioinoculants and graded doses of nitrogen on growth and flower yield of Marigold. Orissa Journal of Horticulture. 1999;27:31-34.
6. Kumar P, Raghava SPS, Misra RL. Effect of Biofertilizers on Growth and Yield of China Aster. Journal of Ornamental Horticulture. 2003;6:85-88.
7. Narendra Kumar, VM Prasad, Netra Pal Yadav. Effect of chemical fertilizers and bio fertilizers on flower yield, tuberos root yield and quality parameter on dahlia (*Dahlia variabilis* L.) cv. Kenya orange. Journal of Pharmacognosy and Phytochemistry. 2019;8(4): 2265-2267.
8. Pandey, Satish & Kumari, Supriya & Singh, Digendra & Singh, Vivek & Prasad, V. Effect of Biofertilizers and Organic Manures on Plant Growth, Flowering and Tuber Production of Dahlia (*Dahlia variabilis* L.) Cv. S.P. Kamala. International Journal of Pure & Applied Bioscience. 2017;5:549-555.

9. Sabah SS. Effect of Different Organic and Inorganic Manure on Flower Yield and Tubers Yield of Dahlia (*Dahlia variabilis* L.) cv. Glory of India as Intercropping with Damask Rose. European Academic Research. 2014;4.
10. Sheergojri GA, Rather ZA, Khan FU, Nazki IJ, Qadriz A. Effect of chemical fertilizers and Bio inoculants on growth and flowering of Dahlia (*Dahlia variabilis*, Desf.) cv. Pink Attraction; 2013.

© 2021 Raghunandan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/70294>