



Role of Oxidative Stress as a Marker of Cardiovascular Risk in Children on Regular Hemodialysis

Muhammad Abrar ^{a*}, Mazhar Nadeem ^a and Sunila Fatima ^b

^a Sahiwal Medical College, Sahiwal, Pakistan.

^b DHQ Hospital, Faisalabad, Pakistan.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i59B34398

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/79686>

Original Research Article

Received 08 October 2021

Accepted 16 December 2021

Published 18 December 2021

ABSTRACT

Introduction: Chronic kidney disease (CKD) is a major public health problem worldwide, and its main consequences include loss of renal function leading to end-stage renal disease (ESRD), increased risk of cardiovascular disease (CVD), significant increase in morbidity and mortality, and a decrease in health-related quality of life.

Aims and Objectives: The basic aim of the study is to analyze the oxidative stress and total antioxidant capacity as a biomarker of cardiovascular risk in those children who are on regular hemodialysis.

Materials and Methods: This cross sectional study was conducted at DHQ hospital, Faisalabad during July 2020 to January 2021. The data were collected from the age of less than 18 years children of both sexes. There were 50 children who was selected for this study. At the time of the study, all the patients were on regular three HD sessions per week. In HD patients, venous blood samples were drawn immediately before and after hemodialysis session. Baseline laboratory investigations were carried out for all patients and controls including complete blood count, serum urea and creatinine, arterial pH, arterial blood gases and infection screening, which included blood and urinary cultures by standard methods.

Results: The data were collected from 50 dialysis patients. The mean age of this study is 15years. We collected all the demographic data of patients. The mean value of Urea is 64.34 ± 2.44 mg/dl). At before-dialysis session, duration of disease positively correlated with TPX ($r = 0.969$, $P < 0.001$),

but, negatively correlated with TAC ($r = -0.469, P < 0.002$). At after-dialysis session, HIF-1 α negatively correlated with each of TPX ($r = -0.529, P < 0.001$) and OSI ($r = -0.459, P < 0.003$); while, OSI positively correlated with TPX ($r = 0.944, P < 0.001$).

Conclusion: It is concluded that HD patients, the clinical and prognostic significance of oxidative status associated with cardiovascular risk factors is very different from the general population. Although a direct causality cannot be inferred from such kind of correlative investigations.

Keywords: Cardiac; risk factors; CKD; prognosis.

1. INTRODUCTION

Chronic kidney disease (CKD) is a noteworthy general health issue around the world, and its fundamental outcomes incorporate loss of renal capacity prompting end-organize renal malady (ESRD), expanded danger of cardiovascular ailment (CVD), critical increment in horribleness and mortality, and a diminishing in health-related personal satisfaction [1,2]. Chronic kidney sickness (CKD) is connected with a truly expanded danger of cardiovascular dismalness and mortality [3]. Different assistant and helpful changes of the cardiovascular structure, for instance endothelial brokenness, vein setting, left ventricular hypertrophy (LVH) and updating of the vessel divider with hyperplasia and calcification occur from the get-go all through CKD (mastermind 2-4 CKD) and add to the undeniable danger of ischemic cardiovascular sickness (CVD) and sudden heart death [4]. While a hindered renal limit can bother "customary" chance variables like hypertension, dyslipidaemia, exacerbation, and oxidative weight, the going with disintegrating of mineral homeostasis and subsequently furthermore bone processing is in all likelihood the key player inciting animated CVD [5]. To highlight the central occupation of mineral assimilation for both, cardiovascular and skeletal genuineness, the term chronic kidney illness mineral bone issue (CKD-MBD) was created starting late [6].

Cardiovascular illness (CVD) is the huge explanation behind dreariness and mortality in patients with end-orchestrate renal ailment (ESRD) on haemodialysis (HD) [7,8]. Since ESRD once in a while results from hypertension and diabetes mellitus, the expanded CVD hazard in these patients has been believed to be the delayed consequence of these essential diseases [9]. Before long, it has been illuminated how ESRD addresses fundamentally a CVD hazard factor self-sufficiently by both hypertension and diabetes mellitus [10].

This incited the wide affirmation that chronic cell hypoxia of the kidney is the last ordinary pathway in the development of CKD provoking inescapable kidney disillusionment [11]. A get-together of elucidation factors, appointed hypoxia inducible interpretation factors, are unequivocally prompted by low tissue oxygen strain and are most likely going to have work in the oxygen-recognizing segment and reparative reaction [12].

1.1 Aims and objectives

The basic aim of the study is to analyze the oxidative stress and total antioxidant capacity as a biomarker of cardiovascular risk in those children who are on regular hemodialysis.

2. METHODS

This cross sectional study was conducted at DHQ hospital, Faisalabad during July 2020 to January 2021. The data were collected from the age of less than 18 years children of both sexes. There were 50 children who was selected for this study. At the season of the examination, every one of the patients were on ordinary three HD sessions for every week; each time for 3– 4 h (all out 12 h week by week) for over 3 months with polysulfone dialyzing films, after creatinine freedom had fallen underneath 8– 12 or potentially pharmacological treatment and diet had demonstrated lacking to control clinical side effects. The mean dialysis term was 2.18 ± 1.36 .

In patients, blood tests were drawn preceding and after hemodialysis session. Pattern lab examinations were done for patients and controls including total blood check, serum urea and creatinine, blood vessel pH, blood vessel blood gases and disease screening included blood and urinary societies by standard strategies.

2.1 Oxidative Status

Oxidative pressure and natural cell reinforcement potential assurance were performed by utilizing

photometric estimation units and a free extreme analyzer framework gave spectrophotometric gadget peruser. The BAP test gives a general proportion of the organic cell reinforcement potential estimating the blood grouping of cancer prevention agents equipped for decreasing the iron from ferric to the ferrous structure (cancer prevention agent hindrance). Results are communicated in $\mu\text{mol/L}$ of the decreased ferric particles.

2.2 Statistical Analysis

Student's t-test was performed to evaluate the differences in roughness between groups. Two-way ANOVA was performed to study the contributions. A chi-square test was used to examine the difference in the distribution of the fracture modes (SPSS 19.0 for Windows, SPSS Inc., USA).

3. RESULTS

The information were gathered from 50 dialysis patients. The mean age of this investigation is 15years. We gathered all the statistic information of patients. The mean estimation of Urea is 64.34 ± 2.44 mg/dl). All the statistic values are introduced in Table 1.

At before-dialysis session, span of malady emphatically associated with TPX ($r = 0.969$, $P < 0.001$), at the same time, adversely connected with TAC ($r = -0.469$, $P < 0.002$). At after-dialysis session, HIF-1 α adversely corresponded with each of TPX ($r = -0.529$, $P < 0.001$) and OSI ($r = -0.459$, $P < 0.003$); while, OSI decidedly associated with TPX ($r = 0.944$, $P < 0.001$).

4. DISCUSSION

Relative hypoxia, as the critical activator of hypoxia-inducible factor, is discernible in chronic kidney malady tissues free of etiology and is thought to result from a mix of fundamental and helpful changes that consolidate; decreased peritubular circulation system related with glomerular harm, hairlike rarefaction, vasoconstriction, luminal narrowing of atherosclerotic vessels [13], expanded oxygen ask for from hyperfiltration and adjusted hypertrophy, confined oxygen dispersal as a result of extracellular framework improvement, and renal shortcoming [14]. Chronically debilitated patients on HD exist in uncommon condition in light of the way that their survival is poor upon medicines which are operational simply 12– 18 h for every week in six hour sessions. This technique subjects these patients to inestimable startling changes in the internal condition, especially fast moves in pH. In this examination, the before-dialysis measurements of plasma lactate, VEGF were out and out higher stood out from healthy controls while, lactate/pyruvate extent was fundamentally higher appeared differently in relation to after-dialysis level [15]. The lactate level would when all is said in done fall with dialysis, at any rate to some degree inferable from this strategy. In like manner plasma level was fundamentally lifted in after-dialysis session appeared differently in relation to healthy controls [16]. Hypoxia is joined by an immense augmentation in blood lactate and outrageous fundamental acidosis as a quick effect of anaerobic assimilation [17]. Beside the quick effects of anaerobic absorption, catecholamine-provoked instigation of cell glycolysis and resulting mix of lactate heightens the expanded fundamental lactate. In such conditions, amassed pyruvate is prepared into lactate [18].

Table 1. Statistic values

Parameters	Mean values
Albumin (g/dl)	6.567 \pm 3.27
Cholesterol, total (mg/dl)	154.8 \pm 4.21
HDL cholesterol (mg/dl)	44.65 \pm 3.21
LDL cholesterol (mg/dl)	79.65 \pm 3.66
C reactive protein (mg/dl)	12.01 \pm 2.11
Uric acid (mg/dl)	5.76 \pm 0.19
Urea (mg/dl)	64.34 \pm 2.44
Iron, total ($\mu\text{g/dl}$)	68.34 \pm 4.81
Transferrin (mg/dl)	149 \pm 5.17
Ferritin (ng/mL)	506.9 \pm 6.21
Protein, total (g/dL)	6.10 \pm 0.12

Table 2. Correlations among the measured parameters in CKD patients at before- and after-dialysis settings

Parameters	Duration	Pyruvate	Lactate/Pyruvate ratio	HIF-1 α	VEGF	TAC	Total peroxides
Pyruvate							
Before dialysis	-0.193(0.234)						
After dialysis	-0.095(0.580)						
HIF-1α							
After dialysis	-0.186(0.249)	0.015(0.929)	0.229(0.156)				
TAC							
Before dialysis	0.265 (0.098)	-0.147(0.366)	-0.142(0.384)	0.068(0.676)	-0.206(0.202)		
After dialysis	0.022 (0.894)	-0.148(0.361)	0.105(0.517)	-0.220(0.173)	-0.147(0.366)		
Total peroxides							
Before dialysis	-0.017(0.918)	-0.194(0.230)	0.128(0.430)	-0.138(0.394)	0.229(0.156)	-0.247(0.124)	
After dialysis	0.283 (0.077)	0.070(0.668)	-0.204(0.207)	-0.529(0.001)	-0.006(0.972)	0.168(0.301)	
Oxidative stress index							
Before dialysis	-0.093(0.566)	0.161(0.322)	0.156(0.337)	-0.172(0.287)	0.245(0.128)	-0.469(0.002)	0.969(0.001)
After dialysis	0.285 (0.075)	0.113(0.489)	-0.240(0.136)	-0.459(0.003)	0.062(0.705)	-0.155(0.340)	0.944(0.001)

Data shown are r value (P< value). HIF-1 α = hypoxia induced factor-1 α , VEGF= vascular endothelial growth factor. TAC = total antioxidant capacity

5. CONCLUSION

It is inferred that HD patients, the clinical and prognostic centrality of oxidative status related with cardiovascular hazard factors is altogether different from the all inclusive community. Despite the fact that an immediate causality can't be construed from such sort of correlative examinations.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors

CONSENT AND ETHICAL APPROVAL

As per university standard guideline, participant consent and ethical approval have been collected and preserved by the authors

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Elshamaa MF, Sabry S, Nabih M, Elghoroury EA, El-Saaied GS, Ismail AAG. "Oxidative stress markers and C-reactive protein in pediatric patients on hemodialysis," *Annals of Nutrition & Metabolism*. 2009;55(4):309–316.
2. Kooman JP, Kotanko P, Schols AMWJ, Shiels PG, Stenvinkel P. Chronic kidney disease and premature ageing," *Nature Reviews Nephrology*. 2014;10(12):732–742.
3. Young PP, Hofling AA, Sands MS. VEGF increases engraftment of bone marrow-derived endothelial progenitor cells (EPCs) into vasculature of newborn murine recipients. *Proc Natl Acad Sci USA*. 2002;99(18):11951–11956.
4. Zhao Q, Ishibashi M, Hiasa K, Tan C, Takeshita A, Egashira K. Essential role of vascular endothelial growth factor in angiotensin II-induced vascular inflammation and remodeling. *Hypertension*. 2004;44(3):264–270.
5. Gunaratnam L, Bonventre JV. HIF in kidney disease and development. *J Am Soc Nephrol*. 2009;20(9):1877–1887.
6. Poellinger L, Johnson RS. HIF-1 and hypoxic response: the plot thickens. *Curr Opin Genet Dev*. 2004;14(1):81–85.
7. Galli F, Piroddi M, Bartolini D, et al. Blood thiol status and erythrocyte glutathione-S-transferase in chronic kidney disease patients on treatment with frequent (daily) hemodialysis," *Free Radical Research*. 2013;48(3):273–281.
8. Stępniewska J, Gołembiewska E, Dołęgowska B, Domański M, Ciechanowski K. Oxidative stress and antioxidative enzyme activities in chronic kidney disease and different types of renal replacement therapy," *Current Protein & Peptide Science*. 2015;16(3):243–248.
9. Harma M, Harma M, Erel O. Measurement of the total antioxidant response in preeclampsia with a novel automated method. *Euro J Obst Gyn Reprod Biol*. 2005;118(1):47–51.
10. Fine LG, Norman JT. Chronic hypoxia as a mechanism of progression of chronic kidney diseases: from hypothesis to novel therapeutics. *Kidney Int*. 2008;74(7):867–872.
11. Yee Koh M, Spivak-Kroizman TR, Powis G. HIF-1 regulation: not so easy come, easy go. *Trends Biochem Sci*. 2008;33(11):526–534.
12. Higgins DF, Kimura K, Bernhardt WM, Shrimanker N, Akai Y, Hohenstein B, Saito Y, Johnson RS, Kretzler M, Cohen CD, Eckardt KU, Iwano M, Haase VH. Hypoxia promotes fibrogenesis in vivo via HIF-1 stimulation of epithelial-to-mesenchymal transition. *J Clin Invest*. 2007;117:3810–3820.
13. Neusser MA, Lindenmeyer MT, Moll AG, Segerer S, Edenhofer I, Sen K, Stiehl DP, Kretzler M, Gröne HJ, Schlöndorff D, Cohen CD. Human nephrosclerosis triggers a hypoxia-related glomerulopathy. *Am J Pathol*. 2010;176(12):594–607.
14. Chiang CK, Tanaka T, Inagi R, Fujita T, Nangaku M. Indoxyl sulfate, a representative uremic toxin, suppresses erythropoietin production in a HIF-dependent manner. *Lab Invest*. 2011;91(11):1564–1571.
15. Stevens LA, Coresh J, Greene T, Levey AS. Assessing kidney function-measured

- and estimated glomerular filtration rate. The New England Journal of Medicine. 2006;354(23):2473–2483.
16. Cornelli U, Terranova R, Luca S, Cornelli M, Alberti A. Bioavailability and antioxidant activity of some food supplements in men and women using the D-Roms test as a marker of oxidative stress. The Journal of Nutrition. 2001;131(12):3208–3211.
 17. La Russa D, Brunelli E, Pellegrino D. Oxidative imbalance and kidney damage in spontaneously hypertensive rats: activation of extrinsic apoptotic pathways. Clinical Science. 2017;131(13):1419–1428.
 18. Matsuyama Y, Terawaki H, Terada T, Era S. Albumin thiol oxidation and serum protein carbonyl formation are progressively enhanced with advancing stages of chronic kidney disease. Clinical and Experimental Nephrology. 2009;13(4): 308–315.

© 2021 Abrar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79686>