



Microbial Quality Assessment, Proximate, Vitamins, and Elemental Analysis of Fermented Milk Vended in Gombe Metropolis, Nigeria

Mohammed Bashir ^{a*}, Mohammed Bilyaminu ^b
and Abdullahi Tawfiq Umar ^c

^a Department of Microbiology, Modibbo Adama University, Yola, Nigeria.

^b Department Science Laboratory Technology, Modibbo Adama University, Yola, Nigeria.

^c Department of Microbiology, Gombe State University, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/SAJRM/2022/v14i3275

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96607>

Original Research Article

Received: 25/10/2022
Accepted: 29/12/2022
Published: 31/12/2022

ABSTRACT

Milk and milk products are highly nutritious substances and they are more prone to microbial attack and spoiled very easily. Fermented milk was analyzed to ascertain the level of contaminants and to also to quantify its nutritional composition. Total aerobic and anaerobic bacteria contaminants as well as coliform bacteria in the fermented milk were determined. The proximate, vitamin and mineral concentration of the fermented milk were also ascertained. The pH, temperature, turbidity and acidity of the fermented milk were found to be 4.38 ± 0.29 , $10.26 \pm 1.7^{\circ}\text{C}$, $6.22 \pm 4.89\text{NTU}$, 0.16 ± 0.04 respectively. The mean aerobic and anaerobic bacteria were $2.23 \times 10^9\text{cfu/mL}$ and $6.15 \times 10^3\text{cfu/mL}$ while the coliform count was $6.10 \pm 2.2\text{MPN/100mL}$. However, the fermented milk analyzed

*Corresponding author: Email: mbashir@mau.edu.ng, mbash1460@gmail.com;

was found to contain an appreciable amount of proximate constituent, vitamins and mineral element. Lack of cooling system in most of the study areas can significantly increase the pH and temperature of the milk and this will create a good environment for microorganisms which when fermented milk stay longer the bacteria will further ferment the milk and produce acid turn the milk sour.

Keywords: Proximate; aerobic and anaerobic count; most probable number; fermented milk; vitamin; elemental analysis.

1. INTRODUCTION

Milk and milk products are important part of traditional food for Africans. Milk and its products are consumed in most of African countries including Nigeria. In Africa the fermented milk and its products were produce using traditional and cultural methods, and fermentation of raw milk is one of the best methods of preserving the nutritional components of milk food [1]. "Most traditional products are prepared without pasteurization by spontaneous fermentation, commonly initiated either by back slopping, i.e., inoculation of fresh milk with the spoiled one, or by repeated use of the same utensils in plastic or natural containers such as milk sacks, calabashes, clay pots, stone jars, and baskets" [1,2].

However, because fermented milk is highly nutritious, it is susceptible to microbial spoilage and different microorganisms such as bacteria and fungi can contaminate and spoiled fermented milk and its products. Microbial contamination of milk usually occurs either from the udder of the animal or within the udder of the animal; it can also occur from the surface of milking tools, and sometimes during handling and storage of milk [3]. Moreover, the health and hygienic conditions of the cow, the environments where the cows are kept and milked, and also the processes and methods used in cleaning and sanitizing the milking equipment as well as those used in milk storage, influence the number of microorganisms in milk [4]. Another important factor that leads to the deterioration of milk and provide good condition for microbial contamination of milk includes temperature and length of storage time of milk. Therefore, this study was designed and conducted to investigate the microbiological qualities of fermented milk as well as the nutritional composition of fermented milk sold within Gombe metropolis of Gombe state Nigeria.

2. MATERIALS AND METHODS

2.1 Description of the Study Area

Gombe is a city in north eastern Nigeria (and a local government area) it is the capital city of Gombe state and has an estimated population of 261,536 people (2006 census). Gombe metropolitan has an area of 52sqkm and situated between latitude $9^{\circ} 30'$ and $12^{\circ} 30'$ N and longitude $8^{\circ} 45'$ and $11^{\circ} 45'$ E of the Greenwich meridian. It has two distinct climates, the dry season (November-March) and the rainy season (April–October) with an average rainfall of 850mm. Gombe is a confluence of economic activities by its position as the meeting point for business people from the surrounding states. The states includes: Borno to the north, Yobe to the east, Taraba and Adamawa to the south and Bauchi to the west [5].

2.2 Source of Sample

Fermented milk was collected from three different markets within Gombe metropolis which includes Gombe main market, Tashan Dukku nono market, and Tashan Shongo nono market.

2.3 Sample Collection

Ninety (90) fresh milk samples were collected for the study, from each of the nono markets; thirty (30) samples were collected. After collection, the samples were transported to the Microbiology Laboratory Gombe State, University, in sterile corked plastic tubes packed in an iced container.

2.4 Physiochemical Analysis

2.4.1 Determination of pH

About 10 mL of the milk sample was measured and dispensed into conical flask and its pH was determined using the pH meter (HI 3220 pH/ORP

meter). The pH meter was standardized using standard buffer of pH 4.0 and 7.0 [6,7].

2.4.2 Determination of temperature

The temperature of the milk samples collected was determined at the time of collecting the samples and the temperature was measured by using thermometer immediately after collecting the samples in the sampling locations [8].

2.4.3 Determination of turbidity

The turbidity of the milk samples was determined according to the FAO manual (2010). In this method, 20mL of the sample was measured and poured into a 50mL conical flask containing 4.0g of ammonium sulphate (NH_4SO_4). The mixture was shaken for 1minute to dissolve the ammonium sulphate (NH_4SO_4) and the mixture was then allowed to stand 5minutes and then filtered into a 100mL conical flask through filter paper. Thereafter, 5mL of the clear filtrate was then transferred into a beaker of boiling water for 5minutes. It was then cooled in cold water. The turbidity was then measured at 500nm by using a UV/ Visible spectrophotometer (GENWAY 63100) [7].

2.4.4 Determination of the total titratable acidity

Using the method described by FAO, [7], in this method about 10mL of the milk sample was measured and dispensed into a conical flask, 3 drops of phenolphthalein was then added. Thereafter, 0.1N NaOH was used to titrate to a noticeable pink colour for end point determination. The acidity will be calculated as lactic acid using the relationship:

$$\% \text{ Lactic Acid} = [\text{Titre value}] \times [0.1\text{N NaOH}] \times [\text{Equivalent Factor}] / \text{Volume of Sample} \times 100$$

Normality of NaOH = 0.1

Volume of Sample = 10ml

Equivalent factor = 0.009. This is because 0.009 gram of lactic acid is found in 1mL of 0.1N lactate.

2.5 Determination of the Total Aerobic and Anaerobic Bacteria

Total aerobic and anaerobic bacterial count was determined by using a pour plate method as described by FAO, [7]. Serial dilution was conducted to obtain 10^{-1} , 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} ,

10^{-6} and 10^{-7} dilutions and 1mL was transferred from each dilution into a duplicate petri dish labeled accordingly. 20-25mL of molten nutrient agar was added to the plates and the plates were allowed to solidify and then incubated at 37°C for 24 hours aerobically and anaerobically. After incubation, plates containing 30-300 colonies were selected and the colonies were counted and recorded. The average was taking and the number obtained was multiplied by inverse of the dilution factor. This gives the number of colony forming units per mL of each sample (cfu/mL).

2.6 Determination of Coliform Bacteria

Coliform count was done using the "Most Probable Number (MPN) technique" also known as the Multiple Tube Fermentation Test (MTFT). In this method, the samples were serially diluted by transferring 1mL of the sample into 9mL of distilled water. The dilutions were 10-fold (10x), 100-fold (100x) and 1000-fold (1000x). Fifteen tubes were groups into fives to make three (3) set. 1mL from each of the dilutions was transferred into first set of 5 tubes containing 9mL of lactose broth and inverted Durham tubes; then 1ml was added to the second set of five tubes and then to the third set. The tubes were incubated at 37°C for 24 – 48hrs. Presumptive positive tubes were recorded for any growth accompanied with gas production in the Durham's tube. Presumption positive tubes were confirmed by sub-culturing two loopful from any tube showing gas production into a tube of brilliant green lactose bile broth containing inverted Durham tubes. The tubes were then incubated at 37°C for 24 – 48hrs; gas production in the Durham's tube confirmed the positive results and the number of confirmed positives tubes were recorded. The Most Probable Number was determined by comparing the number of positives tubes with MPN index table [9].

2.7 Proximate Analysis

The moisture, fat and ash contents of the milk samples were determined using loss in weight after drying to a constant weight as described by AOAC, [6]. The proteins content was determined by using Biuret method. The biuret method is based on the reactions that occur between cupric ions in the reagents and the peptide bonds of the proteins molecules in alkaline solutions to form blue – violet or purple coloured complexes. The absorbance of the colour was measured using a calorimeter at 540 nm [10].

2.8 Mineral Elements Analysis

The presence of mineral elements such as Magnesium (Mg), Manganese (Mn), Calcium (Ca), Potassium (K) and Sodium (Na) in the milk samples was ascertained by using a methods described by Lawal and Adedeji, [11]. Two milliliter (2mL) of each sample was incinerated to a white ash at 550°C in a muffle furnace for 3 hours, allowed to cool and the ash was washed into 250mL beaker with de - ionized water and dissolved with a drop of HCL and filtered using Whatman no 1 filter paper. The volume was made up to 100mL with de-ionized water. The minerals elements were determined using spectrometry method of Atomic Absorption Spectrophotometer (Buck 205) by Buck Scientific. Samples were aspirated and the mean signal response was recorded at each of the element respective wavelength.

2.9 Analysis of Vitamins

The vitamins content of the milk sample was determined using the official methods of the AOAC, [6]. The amount of vitamins such as vitamin A, vitamin C and vitamin E present in the fermented milk were analyzed by using UV/Visible spectrophotometer (GENWAY 63100).

2.10 Statistical Analysis

Analysis of variance protocol was used to determine the level of significance at $p \leq 0.05$ in the mean values of the physiochemical parameters as well as the significance difference in the milk samples collected at different location in the study area using SPSS version 2.0.

3. RESULTS

3.1 Physiochemical Parameters

The results of physiochemical analysis showed that the pH of the milk is acidic and it ranges from 4.19 ± 0.03 to 4.59 ± 0.27 . Samples collected from Tashan Shongo nono market (TSM) has the highest temperature with $10.51 \pm 1.5^\circ\text{C}$; while samples collected from Tashan Dukku nono market (TDM) has the lowest temperature obtained with $9.86 \pm 1.8^\circ\text{C}$. The turbidity of the milk obtained ranges from $1.90 \pm 1.47\text{NTU}$ to $12.07 \pm 2.75\text{NTU}$ while the total titratable acidity

ranges from 0.15 ± 0.03 to 0.18 ± 0.04 as shown in Table 1.

3.2 Total Aerobic and Anaerobic Bacteria Count

The results anaerobic and aerobic bacterial count is described in Table 2. The results revealed that the aerobic bacteria in the fermented milk range from $6.53 \times 10^8\text{cfu/mL}$ to $9.18 \times 10^8\text{cfu/mL}$. Also the anaerobic bacteria count ranges from $3.60 \times 10^3\text{cfu/mL}$ to $1.21 \times 10^3\text{cfu/mL}$.

3.3 Total Coliform Count

The coliform count in the milk samples is presented in Table 3; the results revealed a presence of coliform bacteria which ranges from $5.97 \pm 2.1\text{MPN/100mL}$ to $6.17 \pm 2.4\text{MPN/100mL}$. The overall mean coliform count is $6.10 \pm 2.2\text{MPN/100mL}$.

3.4 Proximate Composition

The milk has high moisture content, which ranges from $8.52 \pm 3.4\text{mg/100mL}$ to $51.53 \pm 12.0\text{mg/100mL}$. The protein contents ranges from $1.58 \pm 0.9\text{mg/100mL}$ to $20.89 \pm 20.1\text{mg/100mL}$ with samples collected from TDM having the lowest protein contents. Fat content is highest in samples collected from TSM with $5.73 \pm 0.6\text{mg/100mL}$ and lowest in samples from GMM with $0.57 \pm 0.4\text{mg/100mL}$. The ash content was found to range from $2.23 \pm 2.6\text{mg/100mL}$ to $2.94 \pm 2.6\text{mg/100mL}$. While the carbohydrate contents of the milk ranges from $40.76 \pm 14.2\text{mg/100mL}$ to $76.88 \pm 9.7\text{mg/100mL}$ as described in Table 4.

3.5 Vitamin Contents

Table 5 described the results of the determination of vitamin contents of the milk samples. The results revealed that values of vitamin A ranges from $47.48 \pm 22.1\text{IU/100mL}$ to $75.23 \pm 25.2\text{IU/100mL}$. Vitamin C content of the samples ranges from $0.59 \pm 0.1\text{mg/100mL}$ to $0.76 \pm 0.2\text{mg/100mL}$. The result also showed that milk samples collected from TSM have the lowest concentration of vitamin C. However, the concentration of vitamin E ranges from $0.58 \pm 0.2\text{mg/100mL}$ to $0.77 \pm 0.3\text{mg/100mL}$.

Table 1. Results of the physiochemical parameters

S/N	Location	Samples Collected	pH	Temperature (°C)	Turbidity (NTU)	Total Titratable Acidity
1	GMM	30	4.59±0.27	10.43±1.8	4.70±2.54	0.18±0.04
2	TDM	30	4.47±0.26	9.86±1.8	1.90±1.47	0.16±0.04
3	TSM	30	4.19±0.34	10.51±1.5	12.07±2.75	0.15±0.03
4	Total	90	4.38±0.29	10.26±1.7	6.22±4.89	0.16±0.04

Key: GMM = Gombe Main Market, TDM = Tashan Dukku Market, TSM = Tashan Shongo Market, NTU = Nephelometric turbidity unit

Table 2. Total Aerobic and Anaerobic Bacterial Count in cfu/mL

S/N	Location	Samples Collected	Aerobic Bacteria count (cfu/mL)	Anaerobic Bacteria Count (cfu/mL)
1	GMM	30	9.18x10 ⁸	1.21x10 ³
2	TDM	30	6.57x10 ⁸	1.34x10 ³
3	TSM	30	6.53x10 ⁸	3.60x10 ³
4	Total	90	2.23x10 ⁹	6.15x10 ³

Key: GMM = Gombe Main Market, TDM = Tashan Dukku Market, TSM = Tashan Shongo Market

Table 3. Total Coliform Count of Fermented Milk MPN/100mL

S/N	Location	Samples Collected	Total Coliform Count
1	GMM	30	6.17±2.4
2	TDM	30	6.17±2.1
3	TSM	30	5.97±2.1
4	Total	90	6.10±2.2

Key: GMM = Gombe Main Market, TDM = Tashan Dukku Market, TSM = Tashan Shongo Market

Table 4. Proximate Composition of Fermented Milk; it is expressed in mg/100mL

S/N	Location	Samples Collected	Moisture	Protein	Fat	Ash	Carbohydrate
1	GMM	30	51.53±12.0	4.28±3.8	0.57±0.4	2.94±2.6	40.76±14.2
2	TDM	30	15.50±8.9	1.58±0.9	3.73±1.3	2.31±2.8	76.88±9.7
3	TSM	30	8.52±3.4	20.89±20.1	5.73±0.6	2.23±2.6	62.54±21.3
4	Total	90	25.18±20.9	8.92±14.5	3.34±2.3	2.49±2.6	60.06±21.6

Key: GMM = Gombe Main Market, TDM = Tashan Dukku Market, TSM = Tashan Shongo Market

Table 5. Vitamin Contents of Fermented Milk

S/N	Location	Samples Collected	Vitamin A (IU/100mL)	Vitamin C (mg/100mL)	Vitamin E (mg/100mL)
1	GMM	30	58.58±21.2	0.76±0.2	0.58±0.21
2	TDM	30	75.23±25.2	0.72±0.2	0.77±0.33
3	TSM	30	47.48±22.1	0.59±0.1	0.62±0.24
4	Total	90	60.43±25.4	0.69±0.2	0.66±0.27

Key: GMM = Gombe Main Market, TDM = Tashan Dukku Market, TSM = Tashan Shongo Market

Table 6. Mineral Constituents of Fermented Milk; it is expressed in mg/100 mL

S/N	Location	Samples	Magnesium	Manganese	Calcium	Potassium	Sodium
1	GMM	30	15.00±20.1	16.37±20.4	217.87±84.9	181.47±15.5	307.97±202.1
2	TDM	30	17.63±20.5	17.67±20.6	225.25±86.8	183.65±16.8	316.67±196.4
3	TSM	30	23.53±20.9	23.13±20.6	251.65±79.1	183.65±16.8	316.83±196.3
4	Total	90	18.72±20.6	19.06±20.5	231.59±83.9	182.92±16.2	313.82±196.1

Key: GMM = Gombe Main Market, TDM = Tashan Dukku Market, TSM = Tashan Shongo Market

3.6 Mineral Constituents

Magnesium, manganese, calcium, potassium and sodium concentration of the samples are reported in Table 6 and the result is expressed in mg/100mL. The magnesium concentration ranges from 15.00 ± 20.1 mg/100mL to 23.53 ± 21.9 mg/100mL. Manganese concentration of the milk is highest in samples collected from Tashan Shongo Nono Market with 23.13 ± 20.6 mg/100mL and lowest in samples collected from Gombe Main Market with 16.37 ± 20.4 mg/100mL. The calcium concentrations ranges from 217.87 ± 84.9 mg/100mL to 251.65 ± 79.1 mg/100mL. The milk has potassium concentration ranging from 181.47 ± 15.5 mg/100mL to 183.65 ± 16.8 mg/100mL. However, concentration of sodium ranges from 307.97 ± 202.1 in GMM to 316.83 ± 196.3 mg/100mL TSM sample.

4. DISCUSSION

The study revealed that there is no significant difference in the pH values between the locations of sample collection. As pH measure the degree of acidity and alkalinity of sample of products, the study showed that all the samples are within the acidic range. The average pH of the fermented milk was found to be 4.4 ± 0.21 . This pH value obtained in this study is closely similar with the pH values by Lawal and Adedeji, [11] on their work "Nutritional and elemental analysis of warankasi (fermented milk product) sold in Lagos metropolis". Analysis of temperature of the milk samples showed that there was significant difference in the temperature of the samples at 5% level of significant between the locations of collecting samples used in this study. The mean of temperature $10.26 \pm 1.7^\circ\text{C}$ obtained in this study is slightly higher than the $7-10^\circ\text{C}$ recommended by FAO, [7]. Higher temperature of 22.83 ± 1.22 was reported by Gemechu et al., [12] in their work on physical and chemical quality of raw cow's milk produced and marketed in Shashemene town, Southern Ethiopia. Lack of cooling system and inefficient use of refrigerator by milk sellers in the study area can increase the temperature of the milk and also increase the number of microbial contaminants [12].

"Studies on the turbidity of fermented milk samples indicated that there was significance difference between locations of collecting samples. The lowest turbidity was found to be 1.90 ± 1.47 NTU. However, the samples fail to satisfy the turbidity standard value of zero (0)

which said that milk is considered sterile when it shows no turbidity" [7]. The mean total titratable acidity of 0.16 ± 0.04 was found in the study. Lawal and Adedeji, [11] reported "higher titratable acidity than those found in this study. Moreover, higher acidity in milk indicated that the milk was kept at room temperature for a longer period of time" [11]. Higher acidity value (0.194 ± 0.006) was also reported by Gemechu et al., [12]. When there is high acidity in milk, the milk can become sour as a result of high accumulation of lactic acid bacteria.

The average mean of aerobic bacteria present in the milk sample was found to be 2.23×10^9 cfu/mL; this value significantly lower than 3.2×10^4 cfu/mL reported by Godic and Slavica, [13]. Nzabuheraheza and Nyiramugwera, [14] also reported 7.21×10^7 cfu/mL in traditionally fermented milk; this result is also higher than the result obtained in this study. The result is also in-line with result reported by Ogbonna, [15] on "microbiological analysis and safety evaluation of nono: A fermented milk product consumed in most parts of Northern Nigeria. This low number in aerobic bacteria count may be due to either high microbiological quality of the milk under process and/or heating process in fermented milk. However, analysis on the anaerobic bacteria count revealed that there was no significance difference in the anaerobic count between the locations of collecting samples. The findings also revealed that the fermented milk samples have an average mean of 6.15×10^3 cfu/mL of anaerobic count". Analysis of the fermented milk samples for assessing the total number of coliform bacteria in the samples by means of coliform count showed that there was a significant difference between the locations of collecting samples. The average mean of coliform bacteria was found to be 6.10 ± 2.2 MPN/100mL. The low coliform count obtained in this study could be as a result of good hygienic method of milking practiced observed by the milk producers.

"Moisture content is a measure of the water content in a product sample" [11]. There was significance difference between the in the moisture content from the sample collected from the three location of collecting samples. The fermented milk has a moisture content of 25.18 ± 20.9 mg/100mL. The result obtained is lower than 68.4% obtained by Lawal and Adedeji, [11]. According to Aworh and Akinniyi, [16] "the moisture content accounts for the textural property of the product sample. The significant of moisture content in milk is that, high moisture

content supports microbial activities and reduces the shelf life of the milk". "Low moisture contents on the other hand, implies low water activities, low water activities causes reduction in microbial growth and the predominant microbial culture consequently increasing the shelf life of the milk samples as a result of low availability of water for microbial growth" [17].

"The protein content of milk is an essential feature of its market value since higher protein content enhances performance of technological transformation" [18]. There was significant difference in protein content between the locations of collecting sample. The results indicated that fermented milk have a mean protein of $8.92 \pm 14.5 \text{mg}/100\text{mL}$. This low protein may be because most of the proteins are destroyed by heat during pasteurization of the milk. The result is lower than the result obtained by Lawal and Adedeji, [11]. These values were higher than that found by Ahmad et al., [19] in raw milk and Gemechu et al., [12].

The fat contents from various locations were found have a significant difference. The average mean obtained is $3.34 \pm 2.3 \text{mg}/100\text{mL}$ and it is lower than those reported by Lawal and Adedeji, [11] and Gemechu et al., [12]. Low fat concentration obtained in this study may result from the fact that during fermentation of milk most fats are removed. Hence the amount of fat present the fermented milk is usually less than those present in the fresh milk.

"The study shows that there was no significant difference in ash contents between the locations of collecting sample. The mean ash content for fermented milk samples was found to be $2.48 \pm 1.9 \text{mg}/100\text{mL}$. This value is lower than the result" reported by Lawal and Adedeji, [11]. The value is higher than the value reported by Ponka et al., [18]; Gemechu et al., [12]. The amount present in a particular sample can determine the amount of mineral element available in that sample.

The carbohydrate content in the sample was obtained by subtracting from 100 the percentage of moisture, protein, fat and ash. The average carbohydrate content is $47.59 \pm 20.9 \text{mg}/100\text{mL}$ which is higher than those reported by Lawal and Adedeji, [11] and higher than those reported by Ajai et al., [17]. Significantly, carbohydrate provides energy to the body [11].

"Vitamin A and E concentration of the samples between the locations of collecting samples were remarkably different. The results of the analysis also showed that the fermented milk samples have an average mean of $60.43 \pm 25.4 \text{IU}/100\text{mL}$. This value is significantly lower than $264.5 \text{IU}/100\text{mL}$ " reported by Yasmin et al., [20]. Vitamin A is important in normal vision, gene expression, growth and immune function by its maintenance of epithelial cell functions [21]. The result also showed that there was significant difference in the vitamin C content between the locations of collecting samples. The results of the analysis indicated that an average mean of $0.69 \pm 0.2 \text{mg}/100\text{mL}$ for fermented milk samples. The concentration of vitamin C obtained in this study is lower than those reported by Yasmin et al., [20]. "Vitamin C is a potent antioxidant that facilitates the transport and uptake of non-heme iron at the mucosa, the reduction of folic acid intermediates and the synthesis of cortisol. It is deficiency of vitamin C fragility to blood capillaries, gum decay and scurvy" [21]. The result showed $0.66 \pm 0.27 \text{mg}/100\text{mL}$ concentration of vitamin E in fermented milk. The result obtained in this study is higher the result reported by Yasmin et al., [20]. "Vitamin E is a powerful antioxidant which helps to protect cells from damage by free radicals and it is vital to the formation and normal function of red blood cell and muscles" [22].

There was no significant difference in the mineral contents of the fermented milk analyzed between the locations of collecting sample. Magnesium concentration of fermented milk has a mean of $18.72 \pm 20.6 \text{mg}/100\text{mL}$. The value obtained was lower than the value reported by Ponka et al., [18] and lower than $12.7-13.7 \text{mg}/100\text{mL}$ reported by Ajai et al., [17]; and higher than $5.12 \text{mg}/100\text{mL}$ reported by Lawal and Adedeji, [11]. There is an overall mean of $251.40 \text{mg}/100\text{mL}$ of calcium content. This value is lower than the value reported by Ajai et al., [17] which ranged between $2000.10 \text{mg}/100\text{mL}$ to $2830.50 \text{mg}/100\text{mL}$. The overall average mean of the potassium (K) content is $159.86 \text{mg}/100\text{mL}$. This value was lower than the value reported by Ajai et al., [17] in Minna Nigeria which ranged between $1065.50 \text{mg}/100\text{mL}$ to $1611.44 \text{mg}/100\text{mL}$. The overall average mean of sodium content is $243.69 \text{mg}/100\text{mL}$. This value is lower than the value $293 \pm 49.39 \text{mg}/100\text{mL}$ reported by Ponka et al., [18] and higher than $2.23 \text{mg}/100\text{mL}$ reported by Lawal and Adedeji, [11].

5. CONCLUSION

The fermented milk analyzed was found to contain high numbers of aerobic and anaerobic bacteria and this high in number may result from lack of cooling system and lack of good storage condition. However, in spite of that, the fermented milk examined contained an appreciable amount of vitamin A, vitamin E and vitamin C. They also contained elements in good quantity of calcium, potassium, sodium, and magnesium but least in manganese although within the recommended values. In terms of proximate composition it was found to contain abundant moisture, protein, carbohydrate, ash and fat.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Beukes EM, Bester BH, Mostert JF. The microbiology of South African traditional fermented milks. *Int. J. Food Microbiol.* 2001;63(3):189–197. Available:[https://doi.org/10.1016/S0168-1605\(00\)00417-7](https://doi.org/10.1016/S0168-1605(00)00417-7)
2. Gonfa A, Foster HA, Holzapfel WH. Field survey and literature review on traditional fermented milk products of Ethiopia. *Int. J. Food Microbiol.* 2001; 68(3):73–186. Available:[https://dx.doi.org/10.1016/S0168-1605\(01\)00492-5](https://dx.doi.org/10.1016/S0168-1605(01)00492-5)
3. Kamana O, Siele C, Liesbeth J, Anastase K, Mieke U. Microbiological Quality and Safety Assessment of the Rwandan Milk and Dairy Chain; International Association for Food Protection. 2014; 77(2):299–307. Available:<http://dx.doi.org/10.4315/0362-028XJFP-13-230>
4. Murphy SC, Boor KJ. Trouble- shooting and causes of high bacteria counts in raw milk. *Dairy Food and Environmental Sanitation.* 2000; 20(8):606-611.
5. Dauda WA, Lawali R. Gombe metropolis road network mapping. *An Integrated Approach: Academic Journal of Interdisciplinary Studies.* 2014; 3(70):102.
6. AOAC. Official methods of food analysis (15th edition). Williams S. (ed.) Association of Official Analytical Chemists, Washington D.C, USA; 2000.
7. FAO. Manual of Food Quality Control12. Quality Assurance in the Food Control Microbiological Laboratory. FAO FIAT Paris; 2010.
8. Teshome G, Fekadu B, Mitiku E. Physical and Chemical Quality of Raw Cow's Milk Produced and Marketed in Shashemene Town, Southern Ethiopia: *Journal of Food and Agricultural Science.* 2015; 5(2): 7–13.
9. Davidson PM, Roth LA, Gambrel-Lenarz SA. Coliform and other Indicator bacteria. In: Wehr, H.M. & Frank, J.F (eds). *Standard Methods for the Examination of Dairy Products.* 17th edition, Washington, American Public Health Association (APHA). 2004; Chapter 7:187-226.
10. Ochei J, Kolhatkar A. *Medical laboratory science theory and practical: Tata McGraw – Hill Publishing Company Limited India;* 2000.
11. Lawal AK, Adedeji OM. Nutritional and elemental analysis of warankasi (fermented milk product) sold in Lagos metropolis, *International Research Journal of Biotechnology.* 2013; 4(6):112-116. Available:<http://www.interestjournals.org/IRJOB>
12. Gemechu T, Fekadu B. and Mitiku E. Physical and Chemical Quality of Raw Cow's Milk Produced and Marketed in Shashemene Town, Southern Ethiopia: *ISABB-Journal of Food and Agricultural Science.* 2015; 5(2):7–13. Available:<https://dx.doi.org/10.5897/ISABB-JFAS2014.0017>
13. Godič TK, Slavica GT. The microbiological quality of raw milk after introducing the two Day's milk collecting system. *Acta Agriculturae Slovenica.* 2008; 92(1):61–74. Available:<http://aas.bf.uni-ij.si>
14. Nzabuheraheza FD, Nyiramugwera AN. Microbiological analysis of traditionally fermented milk sold in Kinigi sector of Musanze district in Rwanda. *African Journal of Food, Agriculture, Nutrition and Development.* 2016; 16(2):10841-10852. Available:<http://dx.doi.org/10.18697/ajfand.74.15105>
15. Ogbonna IO. Microbiological analysis and safety evaluation of nono: A fermented milk product consumed in most parts of Northern Nigeria. *International Journal of Dairy Science.* 2011; 6(3):181-189. Available:<https://dx.doi.org/10.3923/ijds.2011.181.189>

16. Aworh OC, Akinniyi AO. Consumer acceptability and microbial quality of "Soy – warankasi: A Nigerian soft cheese made from a mixture of soy milk and cow milk, Nigerian Food Journal. 1989; 7: 50-57.
17. Ajai AI, Ochigbo SS, Ndamitso MM, Olaoluwajuwon. Proximate and Mineral Compositions of Different Raw Cow's Milks in Minna, European Journal of Applied Engineering and Scientific Research. 2012; 1(1):23-29.
18. Ponka Roger, Beaucher Eric, Fokou Elie, Kansci Germain, Piot Michel, Leonil Joëlle and Gaucheron Frédéric. Composition of Raw Cow Milk and Artisanal Yoghurt Collected in Maroua (Cameroon). African Journal of Biotechnology. 2013; 12(49): 6866-6875.
Available:<http://dx.doi.org/105897/AJB2013.13206>
19. Ahmad S, Gaucher I, Rousseau F, Beaucher E, Piot M, Grongnet FJ, and Gaucheron F. Effects of acidification on physic-chemical characteristics of buffalo milk: A comparison with cow milk. Food Chemistry. 2008; 106(1):11-17.
Available:<https://doi.org/10.1016/j.foodchem.2007.04.021>
20. Yasmin Adeela, Nuzhat Huma, Masood Sadiq Butt, Tahir Zahoor, Muhammad Yasin. Seasonal variation in milk vitamin contents available for processing in Punjab, Pakistan, Journal of the Saudi Society of Agricultural Sciences. 2012; 11:99–105.
Available:<http://dx.doi.org/10.1016/j.jssas.2012.01.002>
21. Achikanu CE, Eze-Stephen PE, Ude CM, Ugwuokolie OC. Determination of the vitamin and mineral composition of common leafy vegetables in South Eastern Nigeria. International Journal of Current Microbiology and Applied Sciences. 2013; 2(11):347-353.
Available:<http://www.ijcmas.com>
22. Lukaski CH. Vitamin and mineral status: Effect on physical performance. Nutrition Research Centre. 2004; 20(7-8):632-644.
Available:<https://dx.doi.org/10.1016/j.nut.2004.04.001>

© 2022 Bashir et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96607>