



Effect of Nitrogen, Phosphorus and Potassium Levels on Yield and Economics of Interspecific Hybrid Cotton (*Gossypium*spp) under Southern Dry Zone of Karnataka

H. M. Honnappa^{1*}, B. G. Shekara¹, Manjanagouda S. Sannagoudar², Lokesh Patil³ and N. Hamsa³

¹*Department of Agronomy, University of Agricultural Sciences, Bengaluru-560065, India.*

²*Division of Grassland and Silvi-Pasture Management, ICAR-Indian Grassland and Fodder Research Institute, Jhansi-284003, India.*

³*Department of Soil Science and Agricultural Chemistry, University of Agricultural Sciences, Bengaluru-560065, India.*

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2019/47402

Editor(s):

(1) Lanzhuang Chen, Professor Laboratory of Plant Biotechnology, Faculty of Environment and Horticulture, Minami Kyushu University, 3764-1, Tateno-cho, Miyokonojo city, Miyazaki, 885-0035, Japan.

Reviewers:

(1) Mr. Habu Saleh Hamisu, National Horticultural Research Institute, Nigeria.
(2) Mehboob-ur-Rahman, National Institute for Biotechnology and Genetic Engineering (NIBGE), Pakistan.
Complete Peer review History: <http://www.sdiarticle3.com/review-history/47402>

Original Research Article

Received 13 November 2018

Accepted 07 February 2019

Published 11 February 2019

ABSTRACT

The field experiment was conducted at ZARS, V. C. Farm, Mandya during Kharif season of 2016 and 2017 in Southern Dry Zone of Karnataka to optimize levels of major nutrients for interspecific hybrid cotton. The experiment on nutrient management was laid out in RCBD with three replications using factorial concept involving nitrogen, phosphorus and potassium levels. The results revealed that application of 150 kg N ha⁻¹ recorded significantly higher seed cotton yield (1857 kg ha⁻¹) and net returns (Rs. 46,198 ha⁻¹) compared to application of 100 kg N ha⁻¹. Among the phosphorus levels, significantly higher seed cotton yield (2046 kg ha⁻¹) and net returns

*Corresponding author: E-mail: honnappahm918@gmail.com;

(Rs. 54,383) were recorded with application of 75 kg P₂O₅ ha⁻¹ over application of 50 kg P₂O₅ ha⁻¹. Among potassium levels, application of 100 kg K₂O ha⁻¹ recorded significantly higher seed cotton yield (1942 kg ha⁻¹) and net returns (Rs. 49,608 ha⁻¹) as compared to application of 50 kg K₂O ha⁻¹.

Keywords: Hybrid cotton; seed cotton yield and economics.

1. INTRODUCTION

Cotton is popularly called as “White Gold” and considered as “King of fibre crops”. It is an important cash crop of global significance, which plays a dominant role in the world agriculture and industrial economy. India is the important grower of cotton on a global scale. It is one of the important cash crops for farming community. It is cultivated in India for fibre and oil purpose. Cotton is an important raw material for the Indian textile industry and contributes at least 65 per cent of its requirements.

Textile industry in India significantly contributes for Indian economy with over 4 million handlooms, 1500 mills, 1.7 million power looms and thousands of garments, hosiery, processing units and provides direct or indirect employment for about 60 million people in the country [1]. In India, cotton is cultivated in an area of 12.70 million ha with a production of 30.50 million bales and productivity of 494 kg ha⁻¹ lint. The productivity is lower in India compared to world (725 kg ha⁻¹ lint). In Karnataka, it is growing in an area of 4.85 lakh hectares with a production of 15.90 lakh bales and productivity of 596 kg ha⁻¹ lint [2].

Among the improved agronomic practices, efficient use of nutrient is one of the ways to get higher cotton productivity. Efficient use of nutrient is highly essential for sustainable agriculture, in the context of declining per capita availability of cultivable land, dwindling natural resources, increasing ecological disturbances and escalating cost of fertilizers. Increasing productivity by bringing new land under cultivation has limited scope makes it obligatory to use of nutrients more efficiently.

Fertilizer management of hybrid cotton is differing distinctly from that of varieties. The cotton varieties have generally poor vegetative and reproductive growth than hybrids. The hybrids have much potential to bear a higher leaf area per plant much earlier to varieties and require relatively higher nutrition in early stages. Similarly, the hybrid cotton plant bears more

fruiting points and larger bolls. The nutritional demands at various stages of growth of hybrid cotton can ultimately decide the seed cotton yield. A careful planning of schedule, quantity of fertilizers and method of application is needed in case of hybrid cotton in order to obtain higher yield. On an average, cotton crop removes N, P and K at the rate of 125, 24 and 37 kg ha⁻¹, respectively for every 10 bales [3]. So, nitrogen, phosphorus and potassium fertilizers use in cotton production remains important. Application of all the three nutrients had effect on lint yield although most of the response was attributed to N (all cultivars) and to some extent P. The results for all quality parameter suggest that K fertilization is a key to better quality [3].

2. MATERIALS AND METHODS

An investigation entitled “Effect of nitrogen, phosphorus and potassium levels on yield and economics of interspecific hybrid cotton (*Gossypium spp*) under Southern Dry Zone of Karnataka” was conducted during *Kharif* season of 2016 and 2017 at Zonal Agricultural Research Station, Vishweshwaraiah Canal Farm, Mandya. The soil of the experimental site is sandy loam in texture and neutral in reaction with a pH of 7.27 and normal in electrical conductivity (0.38 dS m⁻¹). The organic carbon content was 0.46 per cent and low in available N (210.54 kg ha⁻¹), medium in available phosphorus (27.48 kg ha⁻¹) and available potassium (152.20 kg ha⁻¹). The trial was laid out in Randomized Complete Block Design (RCBD) with factorial concept and replicated thrice. There were 18 treatment combinations involving three nitrogen levels (N₁-100 kg ha⁻¹, N₂-150 kg ha⁻¹ and N₃-150 kg ha⁻¹), two phosphorus levels (P₁-50 kg ha⁻¹ and P₂-75 kg ha⁻¹) and three potassium levels (K₁-50 kg ha⁻¹, K₂-75 kg ha⁻¹ and K₃-100 kg ha⁻¹). Farm yard manure was added at the rate of 10 tons ha⁻¹ to the experimental site 15 days prior to sowing and was mixed thoroughly. The fertilizers were applied through soil application as per the treatments. The cotton hybrid (DCH - 32) was shown on 22nd June 2016 and 6th August 2017 with a spacing of 90 cm × 60 cm. The DCH-32 (Jayalakshmi) is world's first interspecific

(*G. hirsutum* × *G. barbadense*) and non Bt-hybrid cotton of an extra-long staple developed and released from Agriculture Research Station, Dharwad, Karnataka in 1981 and was a landmark in heterosis breeding. It has the yield potential of 35 q ha⁻¹ with ginning per cent of 36, basic staple length of 34.0 - 36.0 mm and the duration of the crop is 190 days. It has spinning count of 80, best suited for Karnataka region and usually grown in Southern Karnataka and Maharashtra. The fertilizer nutrient nitrogen, phosphorus and potassium were applied in the form of urea, SSP and MOP, respectively as per the treatments. The 50 per cent of N, 100 per cent P and K was applied as basal dose at the time of sowing and remaining 50 per cent of nitrogen was top dressed in two splits 25 per cent at 50 DAS and 25 per cent at 75 DAS. The

Pendimethalin was sprayed @ 2 ml l⁻¹ as pre-emergence spray at 0.75 kg a.i. ha⁻¹ and later three Hand weeding were carried out at 15, 30 and 50 DAS. Hoeing operation was carried out at 60 DAS to remove the remaining weeds and also to give support to the base of the crop.

The observations on growth parameters were recorded at 60, 90, 120 DAS and at harvest. Seed cotton yield and stalk yield recorded at harvest and calculated based on the yield obtained from each net plot and converted to kg ha⁻¹ and cost benefit ratio was calculated by using gross returns and total cost of cultivation. The data were statistically analyzed by following the method of [4]. The results have been presented and discussed based on pooled data of two years.

Table 1. Seed cotton and stalk yield of hybrid cotton as influenced by different nutrient levels at harvest of the crop

| Treatments | Seed cotton yield (kg ha ⁻¹) | | | Stalk yield (kg ha ⁻¹) | | |
|--|--|--------|--------|------------------------------------|--------|--------|
| | 2016 | 2017 | Pooled | 2016 | 2017 | Pooled |
| Nitrogen (kg ha⁻¹) | | | | | | |
| N ₁ -100 | 1760 | 1659 | 1710 | 3373 | 3050 | 3212 |
| N ₂ -125 | 1852 | 1747 | 1800 | 3551 | 3210 | 3381 |
| N ₃ -150 | 1912 | 1802 | 1857 | 3664 | 3313 | 3488 |
| S.Em± | 39.97 | 37.69 | 34.95 | 76.61 | 69.27 | 65.65 |
| C.D. (p=0.05) | 114.87 | 108.31 | 100.43 | 220.19 | 199.08 | 188.67 |
| Phosphorus (kg ha⁻¹) | | | | | | |
| P ₁ -50 | 1576 | 1486 | 1531 | 3021 | 2732 | 2877 |
| P ₂ -75 | 2106 | 1986 | 2046 | 4037 | 3650 | 3844 |
| S.Em± | 32.63 | 30.77 | 28.53 | 62.55 | 56.56 | 53.60 |
| C.D. (p=0.05) | 93.79 | 88.44 | 82.01 | 179.78 | 162.55 | 154.05 |
| Potassium (kg ha⁻¹) | | | | | | |
| K ₁ -50 | 1640 | 1546 | 1593 | 3143 | 2842 | 2993 |
| K ₂ -75 | 1915 | 1807 | 1861 | 3673 | 3326 | 3499 |
| K ₃ -100 | 1999 | 1885 | 1942 | 3832 | 3465 | 3649 |
| S.Em± | 39.97 | 37.69 | 34.95 | 76.61 | 69.27 | 65.65 |
| C.D. (p=0.05) | 114.87 | 108.31 | 100.43 | 220.19 | 199.08 | 188.67 |
| Interaction (N×P) | | | | | | |
| N ₁ ×P ₁ | 1485 | 1400 | 1442 | 2846 | 2573 | 2709 |
| N ₁ ×P ₂ | 2035 | 1919 | 1977 | 3901 | 3527 | 3714 |
| N ₂ ×P ₁ | 1602 | 1510 | 1556 | 3070 | 2776 | 2923 |
| N ₂ ×P ₂ | 2103 | 1983 | 2043 | 4032 | 3645 | 3838 |
| N ₃ ×P ₁ | 1643 | 1549 | 1596 | 3149 | 2847 | 2998 |
| N ₃ ×P ₂ | 2181 | 2056 | 2118 | 4180 | 3779 | 3979 |
| S.Em± | 56.53 | 53.30 | 49.42 | 108.35 | 97.96 | 92.84 |
| C.D. (p=0.05) | NS | NS | NS | 311.40 | NS | NS |
| Interaction (N×K) | | | | | | |
| N ₁ ×K ₁ | 1533 | 1445 | 1489 | 2939 | 2657 | 2798 |
| N ₁ ×K ₂ | 1843 | 1737 | 1790 | 3532 | 3193 | 3362 |
| N ₁ ×K ₃ | 1904 | 1795 | 1850 | 3650 | 3300 | 3475 |
| N ₂ ×K ₁ | 1670 | 1574 | 1622 | 3200 | 2893 | 3047 |
| N ₂ ×K ₂ | 1896 | 1788 | 1842 | 3634 | 3286 | 3460 |
| N ₂ ×K ₃ | 1992 | 1878 | 1935 | 3818 | 3452 | 3635 |

| Treatments | Seed cotton yield (kg ha ⁻¹) | | | Stalk yield (kg ha ⁻¹) | | |
|---|--|---|--------|--|--------|--------|
| | 2016 | 2017 | Pooled | 2016 | 2017 | Pooled |
| N ₃ ×K ₁ | 1717 | 1619 | 1668 | 3291 | 2976 | 3133 |
| N ₃ ×K ₂ | 1916 | 1806 | 1861 | 3672 | 3320 | 3496 |
| N ₃ ×K ₃ | 2102 | 1982 | 2042 | 4029 | 3643 | 3836 |
| S.Em± | 69.23 | 65.27 | 60.53 | 132.70 | 119.98 | 113.71 |
| C.D. (p=0.05) | NS | NS | NS | NS | NS | NS |
| Interaction (P×K) | | | | | | |
| P ₁ ×K ₁ | 1318 | 1243 | 1281 | 2527 | 2285 | 2406 |
| P ₁ ×K ₂ | 1673 | 1577 | 1625 | 3206 | 2899 | 3052 |
| P ₁ ×K ₃ | 1738 | 1638 | 1688 | 3331 | 3012 | 3171 |
| P ₂ ×K ₁ | 1961 | 1849 | 1905 | 3759 | 3399 | 3579 |
| P ₂ ×K ₂ | 2097 | 1977 | 2037 | 4019 | 3634 | 3827 |
| P ₂ ×K ₃ | 2261 | 2132 | 2196 | 4334 | 3918 | 4126 |
| S.Em± | 56.53 | 53.30 | 49.42 | 108.35 | 97.96 | 92.84 |
| C.D. (p=0.05) | NS | NS | NS | NS | NS | NS |
| Interaction (N×P×K) | | | | | | |
| N ₁ ×P ₁ × K ₁ | 1168 | 1101 | 1135 | 2239 | 2024 | 2132 |
| N ₁ ×P ₁ × K ₂ | 1605 | 1513 | 1559 | 3076 | 2781 | 2929 |
| N ₁ ×P ₁ × K ₃ | 1681 | 1585 | 1633 | 3222 | 2913 | 3067 |
| N ₁ ×P ₂ × K ₁ | 1898 | 1790 | 1844 | 3638 | 3289 | 3464 |
| N ₁ ×P ₂ × K ₂ | 2080 | 1961 | 2021 | 3987 | 3605 | 3796 |
| N ₁ ×P ₂ × K ₃ | 2128 | 2006 | 2067 | 4078 | 3687 | 3883 |
| N ₂ ×P ₁ × K ₁ | 1363 | 1285 | 1324 | 2612 | 2362 | 2487 |
| N ₂ ×P ₁ × K ₂ | 1698 | 1601 | 1649 | 3254 | 2942 | 3098 |
| N ₂ ×P ₁ × K ₃ | 1744 | 1645 | 1694 | 3343 | 3023 | 3183 |
| N ₂ ×P ₂ × K ₁ | 1976 | 1863 | 1920 | 3788 | 3425 | 3607 |
| N ₂ ×P ₂ × K ₂ | 2094 | 1975 | 2034 | 4014 | 3629 | 3822 |
| N ₂ ×P ₂ × K ₃ | 2239 | 2111 | 2175 | 4292 | 3881 | 4087 |
| N ₃ ×P ₁ × K ₁ | 1425 | 1343 | 1384 | 2731 | 2469 | 2600 |
| N ₃ ×P ₁ × K ₂ | 1715 | 1617 | 1666 | 3288 | 2972 | 3130 |
| N ₃ ×P ₁ × K ₃ | 1788 | 1686 | 1737 | 3427 | 3099 | 3263 |
| N ₃ ×P ₂ × K ₁ | 2009 | 1895 | 1952 | 3852 | 3482 | 3667 |
| N ₃ ×P ₂ × K ₂ | 2116 | 1995 | 2056 | 4056 | 3667 | 3862 |
| N ₃ ×P ₂ × K ₃ | 2416 | 2278 | 2347 | 4631 | 4187 | 4409 |
| S.Em± | 97.90 | 92.31 | 85.60 | 187.66 | 169.67 | 160.80 |
| C.D. (p=0.05) | NS | NS | NS | NS | NS | NS |
| <i>Nitrogen levels (N)</i> | | <i>Phosphorus levels (P)</i> | | <i>Potassium levels (K)</i> | | |
| <i>N₁: 100 kg ha⁻¹</i> | | <i>P₁: 50 kg ha⁻¹</i> | | <i>K₁: 50 kg ha⁻¹</i> | | |
| <i>N₂: 125 kg ha⁻¹</i> | | <i>P₂: 75 kg ha⁻¹</i> | | <i>K₂: 75 kg ha⁻¹</i> | | |
| <i>N₃: 150 kg ha⁻¹</i> | | | | <i>K₃: 100 kg ha⁻¹</i> | | |

Table 2. Economics of hybrid cotton as influenced by different nutrient levels

| Treatments | Gross return (Rs. ha ⁻¹) | | | Net returns (Rs. ha ⁻¹) | | | Benefit cost ratio | | |
|--|--------------------------------------|-------|--------|-------------------------------------|-------|--------|--------------------|-------|--------|
| | 2016 | 2017 | Pooled | 2016 | 2017 | Pooled | 2016 | 2017 | Pooled |
| Nitrogen (kg ha⁻¹) | | | | | | | | | |
| N ₁ -100 | 79197 | 74674 | 76935 | 42451 | 37928 | 40190 | 2.15 | 2.02 | 2.09 |
| N ₂ -125 | 83360 | 78598 | 80979 | 46304 | 41542 | 43923 | 2.24 | 2.11 | 2.18 |
| N ₃ -150 | 86020 | 81107 | 83564 | 48655 | 43741 | 46198 | 2.29 | 2.16 | 2.23 |
| S.Em± | 1799 | 1696 | 1747 | 1799 | 1696 | 1747 | 0.048 | 0.045 | 0.047 |
| C.D. (p=0.05) | 5169 | 4874 | 5022 | 5169 | 4874 | 5022 | 0.138 | 0.130 | 0.134 |
| Phosphorus (kg ha⁻¹) | | | | | | | | | |
| P ₁ -50 | 70932 | 66880 | 68906 | 34516 | 30465 | 32491 | 1.94 | 1.83 | 1.89 |
| P ₂ -75 | 94786 | 89372 | 92079 | 57090 | 51676 | 54383 | 2.51 | 2.37 | 2.44 |
| S.Em± | 1469 | 1385 | 1427 | 1469 | 1385 | 1427 | 0.04 | 0.04 | 0.04 |
| C.D. (p=0.05) | 4221 | 3980 | 4100 | 4221 | 3980 | 4100 | 0.11 | 0.11 | 0.11 |

| Treatments | Gross return (Rs. ha ⁻¹) | | | Net returns (Rs. ha ⁻¹) | | | Benefit cost ratio | | |
|--|--------------------------------------|-------|--------|-------------------------------------|-------|--------|--------------------|-------|--------|
| | 2016 | 2017 | Pooled | 2016 | 2017 | Pooled | 2016 | 2017 | Pooled |
| Potassium (kg ha⁻¹) | | | | | | | | | |
| K ₁ -50 | 73794 | 69579 | 71687 | 37477 | 33262 | 35369 | 2.02 | 1.91 | 1.97 |
| K ₂ -75 | 86112 | 81268 | 83690 | 49056 | 44211 | 46633 | 2.32 | 2.19 | 2.26 |
| K ₃ -100 | 89971 | 84832 | 87401 | 52177 | 47038 | 49608 | 2.38 | 2.24 | 2.31 |
| S.Em± | 1799 | 1696 | 1747 | 1799 | 1696 | 1747 | 0.048 | 0.045 | 0.047 |
| C.D. (p=0.05) | 5169 | 4874 | 5022 | 5169 | 4874 | 5022 | 0.138 | 0.130 | 0.134 |
| Interaction (NXP) | | | | | | | | | |
| N ₁ XP ₁ | 66806 | 62990 | 64898 | 30701 | 26885 | 28793 | 1.85 | 1.74 | 1.79 |
| N ₁ XP ₂ | 91588 | 86357 | 88973 | 54202 | 48971 | 51586 | 2.45 | 2.31 | 2.38 |
| N ₂ XP ₁ | 72072 | 67955 | 70014 | 35657 | 31540 | 33598 | 1.98 | 1.86 | 1.92 |
| N ₂ XP ₂ | 94647 | 89241 | 91944 | 56951 | 51545 | 54248 | 2.51 | 2.37 | 2.44 |
| N ₃ XP ₁ | 73917 | 69695 | 71806 | 37192 | 32969 | 35081 | 2.01 | 1.90 | 1.95 |
| N ₃ XP ₂ | 98124 | 92519 | 95321 | 60117 | 54513 | 57315 | 2.58 | 2.43 | 2.51 |
| S.Em± | 2544 | 2398 | 2471 | 2544 | 2398 | 2471 | 0.068 | 0.064 | 0.066 |
| C.D. (p=0.05) | NS | NS | NS | NS | NS | NS | NS | NS | NS |
| Interaction (N×K) | | | | | | | | | |
| N ₁ ×K ₁ | 68987 | 65047 | 67017 | 32980 | 29039 | 31009 | 1.91 | 1.80 | 1.85 |
| N ₁ ×K ₂ | 82914 | 78178 | 80546 | 46167 | 41431 | 43799 | 2.25 | 2.12 | 2.19 |
| N ₁ ×K ₃ | 85691 | 80796 | 83244 | 48207 | 43313 | 45760 | 2.28 | 2.15 | 2.22 |
| N ₂ ×K ₁ | 75128 | 70837 | 72983 | 38811 | 34519 | 36665 | 2.06 | 1.94 | 2.00 |
| N ₂ ×K ₂ | 85320 | 80447 | 82883 | 48263 | 43390 | 45827 | 2.30 | 2.17 | 2.23 |
| N ₂ ×K ₃ | 89631 | 84511 | 87071 | 51837 | 46718 | 49277 | 2.37 | 2.23 | 2.30 |
| N ₃ ×K ₁ | 77267 | 72854 | 75061 | 40640 | 36226 | 38433 | 2.10 | 1.98 | 2.04 |
| N ₃ ×K ₂ | 86203 | 81279 | 83741 | 48836 | 43912 | 46374 | 2.30 | 2.17 | 2.24 |
| N ₃ ×K ₃ | 94591 | 89188 | 91890 | 56488 | 51085 | 53786 | 2.48 | 2.34 | 2.41 |
| S.Em± | 3115 | 2937 | 3026 | 3115 | 2937 | 3026 | 0.083 | 0.079 | 0.081 |
| C.D. (p=0.05) | NS | NS | NS | NS | NS | NS | NS | NS | NS |
| Interaction (P×K) | | | | | | | | | |
| P ₁ ×K ₁ | 59331 | 55942 | 57637 | 23654 | 20265 | 21960 | 1.66 | 1.57 | 1.61 |
| P ₁ ×K ₂ | 75267 | 70968 | 73118 | 38851 | 34552 | 36702 | 2.07 | 1.95 | 2.01 |
| P ₁ ×K ₃ | 78197 | 73730 | 75964 | 41044 | 36577 | 38811 | 2.10 | 1.98 | 2.04 |
| P ₂ ×K ₁ | 88257 | 83216 | 85737 | 51299 | 46258 | 48779 | 2.39 | 2.25 | 2.32 |
| P ₂ ×K ₂ | 94357 | 88968 | 91662 | 56660 | 51271 | 53965 | 2.50 | 2.36 | 2.43 |
| P ₂ ×K ₃ | 101745 | 95933 | 98839 | 63311 | 57499 | 60405 | 2.65 | 2.50 | 2.57 |
| S.Em± | 2544 | 2398 | 2471 | 2544 | 2398 | 2471 | 0.068 | 0.064 | 0.066 |
| C.D. (p=0.05) | NS | NS | NS | NS | NS | NS | NS | NS | NS |
| Interaction (N×P×K) | | | | | | | | | |
| N ₁ ×P ₁ ×K ₁ | 52562 | 49559 | 51061 | 17195 | 14192 | 15694 | 1.49 | 1.40 | 1.44 |
| N ₁ ×P ₁ ×K ₂ | 72220 | 68095 | 70157 | 36114 | 31989 | 34051 | 2.00 | 1.89 | 1.94 |
| N ₁ ×P ₁ ×K ₃ | 75637 | 71317 | 73477 | 38794 | 34474 | 36634 | 2.05 | 1.94 | 1.99 |
| N ₁ ×P ₂ ×K ₁ | 85413 | 80534 | 82973 | 48765 | 43886 | 46325 | 2.33 | 2.20 | 2.26 |
| N ₁ ×P ₂ ×K ₂ | 93608 | 88261 | 90934 | 56221 | 50874 | 53547 | 2.50 | 2.36 | 2.43 |
| N ₁ ×P ₂ ×K ₃ | 95745 | 90276 | 93010 | 57621 | 52152 | 54886 | 2.51 | 2.37 | 2.44 |
| N ₂ ×P ₁ ×K ₁ | 61324 | 57821 | 59573 | 25647 | 22144 | 23896 | 1.72 | 1.62 | 1.67 |
| N ₂ ×P ₁ ×K ₂ | 76401 | 72037 | 74219 | 39985 | 35621 | 37803 | 2.10 | 1.98 | 2.04 |
| N ₂ ×P ₁ ×K ₃ | 78491 | 74008 | 76249 | 41338 | 36855 | 39096 | 2.11 | 1.99 | 2.05 |
| N ₂ ×P ₂ ×K ₁ | 88932 | 83852 | 86392 | 51974 | 46894 | 49434 | 2.41 | 2.27 | 2.34 |
| N ₂ ×P ₂ ×K ₂ | 94239 | 88856 | 91548 | 56542 | 51159 | 53851 | 2.50 | 2.36 | 2.43 |
| N ₂ ×P ₂ ×K ₃ | 100770 | 95014 | 97892 | 62336 | 56580 | 59458 | 2.62 | 2.47 | 2.55 |
| N ₃ ×P ₁ ×K ₁ | 64108 | 60446 | 62277 | 28121 | 24459 | 26290 | 1.78 | 1.68 | 1.73 |
| N ₃ ×P ₁ ×K ₂ | 77181 | 72772 | 74976 | 40455 | 36046 | 38250 | 2.10 | 1.98 | 2.04 |
| N ₃ ×P ₁ ×K ₃ | 80462 | 75866 | 78164 | 42999 | 38403 | 40701 | 2.15 | 2.03 | 2.09 |
| N ₃ ×P ₂ ×K ₁ | 90427 | 85262 | 87844 | 53159 | 47994 | 50576 | 2.43 | 2.29 | 2.36 |
| N ₃ ×P ₂ ×K ₂ | 95225 | 89786 | 92505 | 57218 | 51779 | 54498 | 2.51 | 2.36 | 2.43 |

| Treatments | Gross return (Rs. ha ⁻¹) | | | Net returns (Rs. ha ⁻¹) | | | Benefit cost ratio | | |
|--|--------------------------------------|--------|---|-------------------------------------|-------|---|--------------------|-------|--------|
| | 2016 | 2017 | Pooled | 2016 | 2017 | Pooled | 2016 | 2017 | Pooled |
| N ₃ ×P ₂ ×K ₃ | 108720 | 102510 | 105615 | 69976 | 63766 | 66871 | 2.81 | 2.65 | 2.73 |
| S.Em± | 4406 | 4154 | 4280 | 4406 | 4154 | 4280 | 0.118 | 0.111 | 0.115 |
| C.D. (p=0.05) | NS | NS | NS | NS | NS | NS | NS | NS | NS |
| <i>Nitrogen levels (N)</i> | | | <i>Phosphorus levels (P)</i> | | | <i>Potassium levels (K)</i> | | | |
| <i>N₁: 100 kg ha⁻¹</i> | | | <i>P₁: 50 kg ha⁻¹</i> | | | <i>K₁: 50 kg ha⁻¹</i> | | | |
| <i>N₂: 125 kg ha⁻¹</i> | | | <i>P₂: 75 kg ha⁻¹</i> | | | <i>K₂: 75 kg ha⁻¹</i> | | | |

3. RESULTS AND DISCUSSION

3.1 Seed Cotton Yield

Among nitrogen levels (Table 1), Significantly higher seed cotton yield (1857 kg ha⁻¹) and stalk yield (3488 kg ha⁻¹) was recorded with the application of 150 kg N ha⁻¹ which was found on par with application of 125 kg N ha⁻¹ (1880 and 3381 kg ha⁻¹, respectively), but superior over application of 100 kg N ha⁻¹ (1710 and 3212 kg ha⁻¹, respectively). The ginning percentage and seed index were non-significantly differed due to nitrogen levels. The higher levels of nitrogen increased the seed cotton yield and yield attributing character which was attributed to positive effect of nitrogen it controls new growth, increases nutrient uptake and preventing abscission of squares and bolls thereby retaining higher bolls and also due to increased photosynthetic rate which might have resulted in higher accumulation of metabolites. Similar findings were also observed by several authors [5,6,7]. In phosphorus levels, application of 75 kg P₂O₅ ha⁻¹ were recorded significantly higher seed cotton yield (2046 kg ha⁻¹) and stalk yield (3844 kg ha⁻¹) with the and superior over application of 50 kg P₂O₅ ha⁻¹ (1531 and 2877 kg ha⁻¹, respectively). The higher levels of phosphorus increased the seed cotton yield and yield attributing character may be attributed to increased number of sympodial and monopodial branches would increase the potentiality to bear more leaves which has a positive association between leaf area and leaf area index and in turn it bear more squares since, cotton has auxiliary fruiting habit. These results were in conformity with the findings of several authors [8,9,10]. With respect to potassium levels, significantly higher seed cotton yield (1942 kg ha⁻¹) and stalk yield (3649 kg ha⁻¹) were recorded with the application of 100 kg K₂O ha⁻¹ which was found on par with application of 75 kg K₂O ha⁻¹ (1861 and 3499 kg ha⁻¹, respectively) superior over application of 50 kg K₂O ha⁻¹ (1593 and 2993 kg ha⁻¹, respectively). The higher levels of potassium which increased the seed cotton yield up to 12.97 per cent might be due to potassium

requirement of developing bolls are the major sinks of potassium and uptake is limited during this phase. Similar findings were also observed by [11], Jagadish et al. [12,13]. The interaction between nitrogen and phosphorus levels (N and P), nitrogen and potassium levels (N and K), phosphorus and potassium levels (P and K) and nitrogen, phosphorus and potassium levels (N, P and K) with respect to yield and yield attributes were found non-significant.

4. ECONOMICS

Application of 150 kg N ha⁻¹(Table-2) has recorded higher gross returns (Rs. 83,564 ha⁻¹), net returns (Rs. 46,198 ha⁻¹) and B:C ratio (2.23) followed by application of 125 kg N ha⁻¹ (Rs. 80,979, Rs. 43,923 and 2.18 ha⁻¹, respectively) superior over application of 100 kg N ha⁻¹ (Rs. 76,935, Rs. 40,190, and 2.09 ha⁻¹, respectively). The higher B:C ratio in above treatment was mainly due to the increased seed cotton yield at higher application of nitrogen which in turn increased gross returns. These results are in line with the findings of Gangaiah and Ahlawat [6,10].

Application of 75 kg P₂O₅ ha⁻¹ has recorded higher gross returns (Rs. 92,079 ha⁻¹), net returns (Rs. 54,383 ha⁻¹) and B:C ratio (2.44) superior over application of 50 kg P₂O₅ ha⁻¹ (Rs. 68,906, Rs. 32,491, and 1.89 ha⁻¹, respectively). The higher B:C ratio in above treatment was mainly due to the increased seed cotton yield at higher application of phosphorus and also better utilization of the inputs which in turn increased gross returns. These results are in line with the findings of several authors [14,15,16,17]. Application of 100 kg K₂O ha⁻¹ has recorded higher gross returns (Rs.87,401 ha⁻¹), net returns (Rs. 49,608 ha⁻¹) and B:C ratio (2.31) followed by application of 75 kg K₂O ha⁻¹ (Rs. 83,690, Rs. 46,633 and 2.26 ha⁻¹, respectively) superior over application of 50 kg K₂O ha⁻¹ (Rs. 71,687, Rs. 35,369, and 1.97 ha⁻¹, respectively). The higher B:C ratio in above treatment was mainly due to the increased seed cotton yield at higher application of potassium which in turn increased

gross returns. These results are in line with the findings of several authors [18,10].

The interaction between nitrogen levels and phosphorus levels (N and P), nitrogen levels and potassium levels (N and K), phosphorus levels and potassium levels (P and K) and nitrogen levels, phosphorus levels and potassium levels (N, P and K) with respect to economics were found non-significant (Table 2). However, numerically higher gross returns, net returns and benefit cost ratio were recorded in treatment combination of 150:75:100 kg NPK ha⁻¹ followed by 125:75:100, 100:75:100, 150:75:75 and 125:75:75 kg NPK ha⁻¹ over 100:50:50 kg NPK ha⁻¹.

5. CONCLUSION

Based on the results of the study following agronomic recommendation can be made for cultivation of hybrid cotton under Southern Dry Zone (Zone 6) of Karnataka. The application of nitrogen 125 kg ha⁻¹, phosphorus 75 kg ha⁻¹ and potassium 75 kg ha⁻¹ will be beneficial for getting higher seed cotton yield and net returns in interspecific hybrid cotton.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Jayakumar M, Surendran U, Manickasundram P. Drip fertigation program on growth, crop productivity, water and fertilizer-use efficiency of Bt cotton in Semi-arid Tropical Region of India. *Commun. Soil Sci. Plant. Annu.* 2015;46:293-304.
- Anonymous. All India Co-ordinated Research Project, Annual Report. 2014-15;4.
- Kefyalew G, Roger KT, Kyle WF, Randal KB, William R. Cotton lint yield and quality as affected by applications of N, P, and K fertilizers. *J. Cotton Sci.* 2007;11:12-19.
- Gomez KA, Gomez AA. Stastical procedure for agricultural research. John Wiley and Sons, New Delhi. 1984;680.
- Deepa GS, Aladakatti YR. Effect of nutrient levels and split application of nitrogen and potassium on yield, economics and fibre quality parameters of interspecific Bt cotton. *J. Farm Sci.* 2015;29(2):203-207.
- Bharathi S, Ratna KS, Vamsi KAN, Chenga RV. Effect of nitrogen levels, split application of nitrogen on yield and fibre quality of Bt cotton in vertisols. *J. Cotton Res. Dev.* 2016;30(2):201-204.
- Joao HZ, Ziany NB, Valdinei S, Jose RCB, Jose DCM. Irrigation and nitrogen effects on seed cotton yield, water productivity and yield response factor in semi-arid environment. *Australian J. Crop Sci.* 2016;10(1):118-126.
- Saleem MF, Bilal MF, Awais M, Shahid MQ, Anjum SA. Effect of nitrogen on seed cotton yield and fibre qualities of Bt. cotton cultivars. *J. Animal Plant Sci.* 2010;20(1): 23-27.
- Seema S, Ahlawat IPS, Rana DS. Effect of phosphorus sources and levels on Bt-cotton (*Gossypium hirsutum*) based intercropping systems. *Indian J. Agron.* 2012;57(3):235-240.
- Prakash BH. Response of cotton (*Gossypium spp.*) to nutrient and irrigation levels in Southern Dry Zone of Karnataka. M. Sc. (Agri.) Thesis, Univ. Agric. Sci., Bengaluru, Karnataka; 2017.
- Sakarvadia HL, Polara KB, Parmar KB, Babariya NB, Kunjadia BB. Effect of potassium and zinc on growth, yield, quality parameters and nutrient uptake by Bt cotton. *Asian J. Soil Sci.* 2009;4(1):24-26.
- Mohan D, Raghavaiah DR, Sreenivas G, Sailaja V, Sivasankar A. Response of Bt. cotton hybrid (*Gossypium hirsutum*) to integrated nitrogen management and potassium. *Indian J. Agron.* 2013;58(3): 391-395.
- Ioannis TT, Sergey S, Demetrios B, Theodora M. Effect of potassium fertilization on leaf physiology, fibre yield and quality in cotton (*Gossypium hirsutum* L.) under irrigated Mediterranean conditions. *Field Crops Res.* 2016;193:94-103.
- Shukla UN, Bhale VM, Wanjari SS, Khadse VA. Production potential and economics of cotton (*Gossypium hirsutum*) hybrids under different plant spacing and NPK levels. *Indian J. Agron.* 2014;59(4): 668-671.
- Gadade GD, Gokhale DN, Chavan AS. Performance of *hirsutum* cotton genotypes to different fertilizer levels under high

- density planting system. J. Cotton Res. Dev. 2015;29(1):45-47.
16. Jadhav SG, Chavan DA, Waghmare YM. Effect of plant spacing, growth regulator and nutrient management on yield, quality and economics of Bt cotton. J. Cotton Res. Dev. 2015;29(1):48-52.
17. Pandagale AD, Khargkharate VK, Kadam GL, Rathod SS. Response of Bt cotton (Gossypium hirsutum L.) to varied plant geometry and fertilizer levels under rainfed condition. J. Cotton Res. Dev. 2015;29(2): 260-263.
18. Jagadish K, Yadav MP, Kushwaha SP, Nand K. Effect of different spacing and potassium levels on yield attributes, yield and economics of *hirsutum* cotton. J. Cotton Res. Dev. 2010;24(2):208-209.

© 2019 Honnappa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/47402>