



Enhancement Biochemicals Levels and Improve Histological Tissues of Kidney and Liver Organs for Duckling's Received Aflatoxin B₁ by Using Propolis Extract

**Nahed M. Ayaat^{1*}, E. M. Embaby², Mona M. Abd Elgalil³, Reyad M. El Sharkawy¹,
Mona M. Gouda¹, N. A. Hassan⁴ and Ahmed Shawky⁵**

¹*Department of Botany and Microbiology, Faculty of Science, Benha University, Egypt.*

²*Department of Plant Pathology, National Research Centre, Cairo, Egypt.*

³*Department of Food Toxicology and Contaminants, National Research Centre, Cairo, Egypt.*

⁴*Department of Zoology, Faculty of Science, Benha University, Egypt.*

⁵*Department of Pathology, Faculty of Veterinary Medicine, Benha University, Egypt.*

⁶*Faculty of Science, Benha University, Egypt.*

Authors' contributions

This work was carried out in collaboration between all authors who read and approved the final manuscript.

Article Information

DOI: 10.9734/MRJI/2018/43773

Editor(s):

(1) Dr. Eliton da Silva Vasconcelos, Department of Physiological Sciences, Federal University of São Carlos – UFSCar, Rod. Washington Luiz, São Carlos, SP, Brazil.

Reviewers:

(1) Kemal Çelik, Çanakkale Onsekiz Mart University, Turkey.

(2) Daohong Chen, China.

(3) Andreia Santos, Federal University of Bahia, Brazil.

Complete Peer review History: <http://www.sciencedomain.org/review-history/27920>

Original Research Article

Received 06 August 2018

Accepted 16 November 2018

Published 22 December 2018

ABSTRACT

The aim of the present study was focused to see the effect of propolis as a natural product on duckling's health when exposed to aflatoxin B₁, to enhancement body weight of duckling's received feed contaminated aflatoxin B₁, to enhancement blood function and to improve kidney and liver tissues. The present study show that, birds began to show symptoms of toxicity after 2 weeks. Aflatoxin B₁ alone was found to reduce feed efficiency and poor health can cause by an imbalance of nutrients. Abnormalities symptoms were found with all ducklings received aflatoxin B₁. These

*Corresponding author: E-mail: dr.nahedayyat1@gmail.com;

birds showing slow eating, leg paralysis, slow moving as result of loss functional movement, inflammatory edema of the eyelids (affect eyes), hair loss, and changed in the color when compared with control. All ducklings fed 0.018 ng / ml aflatoxin B₁ diet had significantly ($P < 0.01$ & $P < 0.05$) lower body weight compared with the control group (Un-treated). Whereas, all ducklings received Propolis extracts plus aflatoxin B₁ were enhanced and appear comparable to the control. Propolis extract was found improve significantly ($P < 0.01$ & $P < 0.05$) all body weight gain of all ducklings in the different treatment groups. On the other hand exposure to aflatoxin B₁ can cause several damage to organ systems, increase significantly ($P < 0.01$ & $P < 0.05$) all tested biochemical parameter determined as Urea m mol/L, Creatinine mg/dl, SGPT U/L and SGOT U/L compared with un-treated control. Whereas, all ducklings received Propolis extracts plus aflatoxin B₁ were improve all these parameters. In the affected birds metabolic changes lead to enlargement of liver, kidney and spleen as well as decrease in the size of bursa of fabricus as a result, liver is greatly enlarged, yellow and friable (easily broken), fat accumulates inside the cell of liver as clear vacuoles. Data show that enlargement in kidney weight in contaminated groups . Whereas, all ducklings received Propolis extracts plus aflatoxin B₁ were improved and appear comparable to the control. Also, microscopic examination of liver and kidney tissues showing, changes of histopathological tissues with all ducklings received aflatoxin B₁. Whereas, all ducklings received Propolis extracts plus aflatoxin B₁ were improved and appear comparable to the control.

Keywords: Feed; *Aspergillus flavus*; ducklings; kidney; liver.

1. INTRODUCTION

Feed is the single most important input in increasing chicken and fish culture production and profits. Information about fungi associated with food and feeds is important in assessing the risk of mycotoxin contamination. Feeds are frequently contaminated simultaneously by several moulds. Mould contamination of food and the environment has a potential to produce fungal toxins (mycotoxins) that are harmful to chickens, turkeys, ducks, fish etc. According to WHO, about 25% and 40% of the world's food is contaminated by mycotoxins. These toxins pose serious health concerns to animals as well as human beings [1,2]. The consumption of these mycotoxin-contaminated feedstuffs by animals leads to an adverse effect on animal health and the effects are more serious in monogastric animals depending on the species and the susceptibility to toxins within the species [3]. The carryover of toxins from animal food may have severe consequences on human health. The economic impact of lowered productivity reduced weight gain, reduced feed efficiency, damage to body organs, interference in reproduction is many times greater than that of immediate mortality and morbidity [4]. Mycotoxins in food and feedstuffs affect both the organoleptic characteristics and the nutritive value of feed, leading to the risk of toxicosis. However, the biological effects of mycotoxins depend on the amount ingested by the host, the age, sex and strain of the animal, varieties of occurring toxins, time of exposure to mycotoxins, animal sensitivity and its condition. Mycotoxins can

induce health problems that are specific to each toxin or suppress the immunity power of animals, favouring infections. This is the major reason for the difficulty of diagnosing mycotoxicoses [5]. Commercial feedstuffs are an important component in modern animal husbandry, but there is no information available about fungal contaminations [6]. Since feed accounts for about 50-60% of the variable costs of production, feed quality is crucial to the success of farming operations. Major problems that may result from low-quality feeds are poor appetite, slow growth, high feed conversion ratio, and low survival. These usually develop as a result of problems on quality of raw materials, feed formulation, processing technology, storage, and feed manage-merit. The most serious problems on feed quality are those involving rancidity, aflatoxin contamination, and nutrient loss [7]. Aflatoxicosis represents one of the serious disease of poultry, livestock and other animals. the cause of this disease in poultry and other food-producing animals has been attributed to the ingestion of various feeds contaminated with *Aspergillus flavus*. This toxegenic fungus is known to produce a group of extremely toxic metabolites, of which aflatoxin B₁ (AFB₁) is most potent. Avian species especially chickens; Gosling, duckling and turkey poult are most susceptible to AFB₁ toxicity. The incidence of hepatocellular tumours. Particularly in duckling, is considered to be one of the serious consequences of aflatoxicosis [8]. Propolis is a resinous substance collected by worker bees (*Apis mellifera*) from the bark of trees and leaves of plants. Propolis shows a complex chemical

composition. Propolis shows pharmacological activities, such as antifungal, antibacterial, anticancer or anti-inflammatory to name a few, among other activities, have attracted the researchers' interest [9,10,11].

Because aflatoxigenic fungi and aflatoxins are the most spread, most dangerous and the most effects on health and economics, the aim of the present study was focused to see the effect of propolis as a natural product on duckling's health when exposed to aflatoxin B₁, to enhancement body weight of duckling's received feed contaminated aflatoxin B₁, to enhancement blood function and to improve kidney and liver tissues.

2. MATERIALS AND METHODS

2.1 Fungal Producing Toxin

Aspergillus flavus was propagated as a pure culture in 100 ml yeast extract sucrose (YES) broth containing 2% yeast extract and 20% sucrose/ litre distilled water according to [12]. One disc 5 mm of *A. flavus* was inoculated into each flask (250 ml) having 100 ml of sterile Y.E.S with 2, 5, 10 and 20% of 'methanol extract of propolis' (MEP). Another flask free 'methanol extract of propolis' (MEP) was used as a control. Three replicates for each treatment. All treatments were incubated at 26±2°C. in the dark for 14 days.

2.2 Preparation of Propolis Extract

The hand-collected propolis was stored in a brown bottle and away from exposure to light until further processing. Propolis extracts were prepared as described by Özdemir et al. [13], propolis was prepared by adding 100 g of the collected propolis to 900 mL of 70% methanol to give 10% methanolic extract of propolis (MEP) which extracted and heating for evaporating methanol (at 50°C for 5 hours) and agitating. Water was then added. To optimise purification, centrifugation at high speeds (4,000 rpm) was proposed. All Samples were centrifuged for 25 minutes. The supernatant was stored overnight at ambient temperatures. The supernatant was further filtered through filter paper (Whatman no. 1) and stored at ambient temperatures in a bottle according to [14]. The final solution was termed 'methanol extract of propolis' (MEP) to produce a final solution from various propolis samples [15]. Kept at 4 °C in dark storage until use.

2.3 AFB₁-Contaminated Feed and Propolis Treatments

Upon analysis, the AFB₁-contaminated feed contained the following aflatoxins 0.018 µg/ml of AFB₁ with 2, 5, 10 and 20% of 'methanol extract of propolis' (MEP). AFB₁ was under the detection limit [16].

2.4 The Toxicity of Aflatoxins and Its Control

2.4.1 Experimental design, Birds, the toxicity of *Aspergillus flavus* toxins on body weight and it's improved

A total of 24 ducklings (7-day-old av. 417 g) (mixed sex) were purchased from Fac. of Veterinary Med, Benha Univ. Ducks were weighed and randomly allotted to 6 dietary treatments. The experimental groups were described in Table 1 according to Mehrim and Salem [17]. The experimental diets were formulated based on the [18] recommendations for 2 wk duck starter diets standard. Ducks were housed on a commercial farm. The temperature was maintained at 34°C to the end of the experiment. The overhead light was provided continuously for the entire period of the experiment. Ducklings were randomly divided into 6 treatment groups of ducks and there were 4 replicates for each treatment, then treated for two weeks and fed according to the indicated experimental diets as follows: group 1, received Basal Diets (BD) plus 0.018 ng/ml AFB₁ plus 2% of propolis' (MEP); group 2, treated orally with BD + 0.018 ng / ml AFB₁ +5% of MEP; group 3, treated orally with BD + 0.018 ng / ml AFB₁ +10% of MEP; group 4, received BD + 0.018 ng / ml AFB₁ +20% of MEP; group 5, treated orally with BD + 0.018 ng / ml AFB₁ (as a positive control); group 6, control which received Basal Diets (BD) free from AFB₁ (as a negative control). The ducklings were raised in battery cages, and warm air was used for brooding ducklings during the trial period. The ducks were observed daily for signs of toxicity. Body weights (BW) and growth uniformity, which was expressed as CV of BW were recorded at zero time, and at the end of the feeding experiment after 2 weeks of the experimental period. Mortalities and health status were observed and recorded daily throughout the entire experimental period as described by Wan et al. [19].

Table 1. Description of the experimental groups

Experimental group	Details
G1	Basal Diets (BD) + 0.018 ng / ml AFB ₁ +2% of propolis' (MEP)
G2	BD + 0.018 ng / ml AFB ₁ +5% of MEP
G3	BD + 0.018 ng / ml AFB ₁ +10% of MEP
G4	BD + 0.018 ng / ml AFB ₁ +20% of MEP
G5	BD + 0.018 ng / ml AFB ₁ (as a positive control) free MEP
G6	Basal Diets (BD) free from AFB ₁ (as a negative control)

2.4.2 Blood samples

At the end of the experiment, birds fasted for 12 h. (All ducks were not fed on the day of the sacrifice). After the experimental period (14 days), a 5-mL blood sample was collected from a jugular vein in a test tube without anticoagulant (using 1 ml syringes) for biochemical analysis. Blood samples were incubated at 37°C for 2 h, centrifuged at 1,500 × g for 10 min, and serum was separated and stored in 1.5-mL centrifuge tubes at -20°C until analysis. Blood samples were stored frozen at -20°C. Blood samples obtained from treated and untreated ducklings were analysis according to [19].

2.4.2 Biochemical studied (Serum analysis)

All the biochemical determinations were carried out using commercial kits according to the kits manufactures unless explained as follow:-

2.4.2.1 Determination of urea

Determination of urea level in serum by using the enzymatic colourimetric method of Barham and Trinder [20] by using kit obtained from BioMérieux SA (France).

2.4.2.2 Determination of Creatinine

Creatinine was determined in serum using commercial kits purchased from Stanbio Laboratory, Inc. (San Antonio, Texas, USA) according to the methods of Bartels et al. [21].

2.5 Histopathological Studies

After bleeding, birds were killed by cervical dislocation for organ weight calculation, including liver and kidney, and their relative weights were calculated as (organ weight/ live weight) × 100 [19]. Livers and Kidney specimens from all ducks were dissected immediately after death and fixed in 10% neutral-buffered formal saline for 72 h. All the specimens were washed in tap water for half an hour and then dehydrated in ascending

grades of alcohol, cleared in xylene and embedded in paraffin. Serial sections of 6 µm thick were cut and stained with Haematoxylin and eosin for histopathological investigation according to Drury and Wallington [22], Saddiq and Kalifa [23] then, examined by light microscopy [22,23,24]. Images were captured and processed using Adobe Photoshop version 8.0.

2.6 Statistical Analysis

Obtained data were subjected to statistical analysis of variance according to Snedecor and Cochran [25], and means separation were done according to Duncan [26].

3. RESULTS AND DISCUSSION

3.1 Enhancement of Duckling Health Received Aflatoxin B1

The present study shows that control group (free aflatoxin B₁) was in normal ducklings (Healthy) as showing in Fig. 1.a. Abnormalities symptoms were found with all ducklings received aflatoxin B₁. Data also show that aflatoxin B₁ alone was found to reduce feed efficiency and poor health can be caused by an imbalance of nutrients either by deficiency. All ducklings fed 0.018 ng/100 g feed aflatoxin B1 diet had lower body weight compared with the control group (Untreated). The birds began to show symptoms of toxicity after 2 weeks, external examination of these birds showing birds hazeled together indicating abnormal behaviour, slow eating, slow movement, loss of functional movement, hair loss and change in the colour when compared with control Fig. 1a-b.). Gimeno [27] found that affected ducklings displayed a delay in development, hyperkeratosis of the cornea and the oral mucosa, malformation and bone fragility, leg paralysis, inflammatory oedema of the eyelids, dermatitis, and scarce feathering. The ducklings suffered massive a vitaminosis and deficiency in calcium, phosphorus and manganese absorption. Gimeno [27], Bibu [28]

stated that in the field, animals experiencing a mycotoxicosis may exhibit a few or many of a variety of symptoms, including digestive disorders, reduced feed consumption, unthriftiness, rough hair coat or abnormal feathering, undernourished appearance, low production, poor production efficiency, impaired reproduction and/or a mixed infectious disease profile. Mycotoxins can increase the incidence of disease and reduce production efficiency. Some of the symptoms observed with a mycotoxicosis may, therefore, be secondary, resulting from an opportunistic disease, present because of mycotoxin-induced immune suppression. Immunotoxic effects of mycotoxins are reviewed. Mariana et al. [29] concluded that the toxegenic *Aspergillus flavus* fungus is known to produce a group of extremely toxic metabolites, of which aflatoxin B₁ (AFB₁) is most potent. Avian species especially chickens; Gosling, duckling and turkey poults are most susceptible to AFB₁ toxicity. The incidence of hepatocellular tumours. Particularly in duckling, is considered to be one of the serious consequences of aflatoxicosis. Wan et al. [19] found that bill colour of ducks receiving contaminated corn was discoloured in contrast to the yellow colour of the ducks receiving uncontaminated corn. Gauthier and Ludlow [30] concluded that moulds are capable of reducing the nutritional value of feedstuff as well as elaborating several mycotoxins. Mycotoxin-contaminated feed has adverse effects on animal health and productivity. Regarding nutritional quality, lipids, proteins, and minerals are of essential importance for the proper development and growth of farm animals. Wu et al. [31] stated that *Aspergillus* causes different forms of aspergillosis. The most common form of *Aspergillus* mould infection is brooder

pneumonia, a lung and air-sac disease of chicks. less – common forms of aspergillosis affect eyes, skin, brain, or bones. Chicks affected by brooder pneumonia gasp, lose their appetite, and look sleepy. The disease does n't spread from chick to chick, but the mould can infect many chicks in a group at once, and up to half may die from the infection.

3.2 Improve Body Weight Gain of Duckling Exposed Aflatoxin B₁

Effect of aflatoxin B₁ produced by *Aspergillus flavus* and detoxification of their effects by using a natural product as Propolis extract on feed intake and changes in body weight gain of ducklings in the different treatment groups were recorded in Table 2. The presented data show that aflatoxin B₁ alone was found to reduce feed efficiency and poor health can be caused by an imbalance of nutrients either by deficiency. All ducklings fed 0.018 ng /100 g aflatoxin B₁ diet had significantly (P < 0.01 & P < 0.05) lower body weight compared with the control group (Un-treated). Visual examination revealed that bill colour of duckling receiving 0.018 ng/100 g aflatoxins B₁ contaminated diet was discoloured. Whereas, all ducklings received Propolis extracts in combination plus aflatoxin B₁ were improved and appeared comparable to the control. Propolis extract was found significantly (P < 0.01 & P < 0.05) improve all body weight gain of all ducklings in the different treatment groups. Also, data show that significantly (P < 0.01 & P < 0.05) enhanced body weight gain of all ducklings with increasing the concentration of Propolis extract. The average of body weight gain in control group was 770 g then decreased to 631 g which loosed 139 g equal 18.1 reduction percent with the first



Fig. (1.a). Control duckling group showing normal symptoms of ducklings (Healthy); Fig. (1.b) Duckling with serious aflatoxicosis problem, Aflatoxin B₁ alone showing abnormality symptoms, slow moving and was found to decrease the seize and body weight gain (weight loss), leg abnormalities (leg weakness) and lethargy, appearance are common symptoms of aflatoxin poisoning

group which received feed contaminated *A. flavus* toxin (aflatoxin B₁) and 2% Propolis. While ducklings of the second group which received feed contaminated with *A. flavus* toxin (aflatoxin B₁) and 5% Propolis loosed 114 g equal 19.1 % reduction. Group 3 was found to reduce from 770 g to 671 g which loose 99 g equal 12.9% reduction when treated with 10% Propolis extract, group 4 loosed 87 g and gave 11.3% reduction when treated with 20% Propolis extract. Group 5 free Propolis extract which received feed contaminated with *A. flavus* toxin (aflatoxin B₁ only) was found to decrease from 770 g to 500 g and gave 95.59% reduction. [32,33] said the economic impact of lowered productivity, reduced weight gain, reduced feed efficiency, damage to body organs, interference in reproduction is many times greater than that of immediate mortality and morbidity. [34] Concluded that in animals, adverse effects of AF also include a reduction in growth rate and feed efficiency, decreased egg production and hatchability, and increased susceptibility to disease. In addition, residues of AF from animals can appear in edible animal products for human consumption, which raises public health concerns. [35,36] found that in chicken, B₁ is metabolised into M₁ and B₂ in the liver and NADP linked enzyme system reduces B₁ and B₂ to cyclopentanol and aflatoxin. Aflatoxin B₁ impairs all-important production parameters in poultry including weight gain, feed intake, feed conversion efficiency, pigmentation, processing yield, egg production, and male and female reproductive performance. Wan et al. [19] said all ducklings fed 100% M diet had lower growth performance ($P < 0.05$) ADG compared with 0%. Chen et al. [36], Mabbett [37], Radziejewska et al. [38] reported that dietary AF exposure reduces weight gain and feed intake, and worsens feed efficiency. The response of animals to AF-contaminated feed depends on the AF concentration, animal species, and age and

sex. The reduced growth rate because of AF ingestion in the diet is primarily due to the reduction in feed intake. Generally, 0.95 mg/kg AF in the diet reduces weight gain by 11 percent because of, in part, reduced feed intake and metabolic inefficiencies from liver and GIT damage. Radziejewska et al. [38], Greco et al. [39], Chang et al. [40] found that most mycotoxicoses of poultry are caused by an intake of low concentration of contaminants over a long whit the typical chronic symptoms of poor growth, poor feed efficiency, and suboptimal production. Ingestion of higher concentration, however, leads to acute clinical symptoms associated with specific vital organs, the immune system, and other aspects of avian physiology as well as mortality. Akande et al. [41] reported that aflatoxins contained in the mould-exposed diet significantly reduced daily weight gain and feed intake of male ducks. Compared with normal-diet group, presence of AFs in the diet significantly reduced weights of meat male ducks at 7, 14, 21, 28, and 35 days of age, significantly reduced daily weight gain and daily feed intake at different periods (1e7 days of age, 15e21 days of age, 22e28 days of age, 29e35 days of age, and significantly increased their F:G at different ages (except for days 8e14; $P < 0.05$).

3.3 Improve Biochemical Function (Improve Blood Function)

Effect of aflatoxin B₁ on blood function as result of biochemical analysis presented that, aflatoxin B₁ can cause damage to organ systems, increase significantly ($P < 0.01$ & $P < 0.05$) all tested biochemical parameter determined as Urea m mol/L, Creatinine mg/dl, SGPT U/L and SGOT U/L compared with un-treated control group 6 (Free aflatoxin B₁) as shown in Table 3. The effects of aflatoxin B₁ for two weeks on serum biochemical parameters of ducklings

Table 2. Effect of aflatoxin B1 on body weight gain of ducklings and their improvement after two weeks by Propolis

Groups	BW (g)		L. (g)	%L	%R
	Zero time	After 14 days			
G1	400.00	631.00	139	81.9	18.1
G2	406.00	622.67	114	80.9	19.1
G3	404.00	671.00	99	87.1	12.9
G4	417.00	683.00	87	88.7	11.3
G5	466.00	500.00	34.00	4.41	95.59
G6 (control)	410.00	770.00			
L.S.D@1%	2.164	24.193			
L.S.D@5%	1.488	16.629			

*BW=Body Weight; L=Loss; %L = Loss percent; %R=Reduction

compared with control group indicated that the control group was in the normal limit, but were found to the differentiation between biochemical parameters i.e. urea, Creatinine, SGPT U/L and SGOT U/L when ducklings received aflatoxin B₁ for two weeks. Higher significant ($P < 0.01$ & $P < 0.05$) was found in between aflatoxin B₁ alone group 5 and the control group 6 (Free aflatoxin B₁). On the other hand analysis of liver function resulted that, significantly ($P < 0.01$ & $P < 0.05$) increase SGPT from 34.33 U/L to 56.67 U/L with 65.1% reduction for group 1 (ducklings received aflatoxin B₁ plus 2% propolis extract), group 2 received aflatoxin B₁ plus 5% propolis extract record 52.67 U/L equal 53.4% reduction, group 3 received aflatoxin B₁ plus 10% propolis extract gave 49.00 U/L and 42.7% reduction, group 4 received aflatoxin B₁ plus 20% propolis extract gave 41.0 U/L and 19.4% reduction while, group 5 received aflatoxin B₁ contaminated diet feed had increase ($P < 0.01$ & $P < 0.05$) serum SGPT U/L and SGOT U/L concentrations as well as increase ($P < 0.01$ & $P < 0.05$) Urea m mol/L, and Creatinine mg/dl concentration compared with concentrations in those fed diet free aflatoxin B₁. it also showed that an increase of SGOT U/L increased from 63.67 to 79.00 in control (free from aflatoxin) with 24.1% reduction while, group 2 received aflatoxin B₁ plus 5% propolis extract record 76.33 equal 19.9% reduction group 3 received aflatoxin B₁ plus 10% propolis extract gave 72.00 U/L and 13.1 reduction, group 4 received aflatoxin B₁ plus 20% propolis extract gave 65.33 and 2.6% reduction while group 5 received aflatoxin B₁ only (free propolis extract) gave 84.00 equal 31.9% reduction. On the other hand analysis of kidney function resulted that, significantly ($P < 0.01$ & $P < 0.05$) increase creatinine from 0.45 to 0.55 with 22.2% reduction for group 1 (ducklings received aflatoxin B₁ plus 2% propolis extract), group 2 received aflatoxin B₁ plus 5% propolis extract record 0.53 equal 17.8% reduction, group 3 received aflatoxin B₁ plus 10% propolis extract gave 0.52 and 15.6 reduction, group 4 received aflatoxin B₁ plus 20% propolis extract gave 0.45 and 00% reduction while, group 5 received aflatoxin B₁ only (free propolis extract) gave 0.56 equal 24.4% reduction. the table also showed Urea conc which increase from 14.67 to 18.7 with 27.5% reduction for group 1 (ducklings received aflatoxin B₁ plus 2% propolis extract), group 2 received aflatoxin B₁ plus 5% propolis extract record 17.3 equal 17.9% reduction, group 3 received aflatoxin B₁ plus 10% propolis extract gave 16.0 and 9.1 reduction, group 4 received aflatoxin B₁ plus 20% propolis extract gave 15.0

and 2.2% reduction while, group 5 received aflatoxin B₁ only (free propolis extract) gave 19.0 equal 29.5% reduction. Sultana and Hanif [42], Grenier and Applegate [43] reported that mycotoxins can cause damage to organ systems, reduce production and reproduction, and increase diseases by reducing immunity. Some mycotoxins are carcinogens, some target liver, kidney, digestive tract or the reproductive system. Bintvihok [33] stated that, in contrast, recent literature reported adverse effects at concentrations as low as 0.02 mg/kg. A plausible explanation of these differences between earlier and more recent reports could be that modern broilers have more efficient nutrient conversion demanding faster hepatic metabolism, which in turn results in a higher metabolism of AFB₁. Anjum [34], Radziejewska et al. [38] reported that the lower tolerance of ducks also could be explained by a lower activity of hepatic enzymes responsible for cellular detoxification and excretion of a variety of toxic substances. In contrast, recent literature reported adverse effects at concentrations as low as 0.02 mg/kg. A plausible explanation of these differences between earlier and more recent reports could be that modern broilers have more efficient nutrient conversion demanding faster hepatic metabolism, which in turn results in a higher metabolism of AFB₁. Ortatatli and Oguz [44], Verma et al. [45], Radziejewska et al. [38] concluded that enzyme activities are modulated following AF consumption. An increased release of enzymes from the pancreas to the intestinal tract might be a consequence of pancreatic damage. Very low doses of AFB₁ (0.02 and 0.04 mg/kg) have reduced the apparent digestibility of crude protein by 8-13 percent in ducks. Similarly, it has been suggested that dietary AF increases the amino acid requirements, and it appears to negatively impact ducks more than chickens.

3.4 Effect of Aflatoxin B₁ Exposure on Intestinal Morphology and its Improved

Effect of aflatoxin B₁ exposure on intestinal morphology consumption of aflatoxin B₁ cause a few or many of a variety of symptoms, including: opportunistic disease, present because of mycotoxin-induced immune suppression, however leads to acute clinical symptoms associated with specific vital organs, Liver as a key player of AF toxicity and sensitivity within Poultry. Liver damage occurred. In the affected bird's metabolic changes lead to enlargement of liver, kidney and spleen as well as a decrease in

Table 3. Improve biochemical changes (improve kidney and liver functions)

Groups	Blood functions (Enzymatic analysis)							
	Liver functions				Kidney functions			
	SGPT U/L	R%	SGOT U/L	R%	Creatinine mg/dl	R%	Urea m mol/L	R%
G1	56.67	65.1	79.00	24.1	0.55	22.2	18.7	27.5
G2	52.67	53.4	76.33	19.9	0.53	17.8	17.3	17.9
G3	49.00	42.7	72.00	13.1	0.52	15.6	16.0	9.1
G4	41.00	19.4	65.33	2.6	0.45	0	15.0	2.2
G5	60.00	74.8	84.00	31.9	0.56	24.4	19.0	29.5
G6	34.33		63.67		0.45		14.67	
(control)								
L.S.D@1%	13.327		20.521		0.048		2.485	
L.S.D@5%	9.160		14.105		0.033		1.533	

* SGPT = Serum glutamic pyruvic transaminase; SGOT= Serum glutamic oxaloacetic transaminase

the size of bursa of fabrics as shown in Figs. 2-4 as a result, the liver is greatly enlarged, yellow and friable (easily broken) compared with the control group (free aflatoxin B₁). Fat accumulates inside the cell of the liver as clear vacuoles compared with control group (free aflatoxin B₁). While duckling received aflatoxin B₁ plus propolis extract was enhanced. Continuous enhancement with increasing propolis extract comparable to control. Similar results were obtained by Andretta et al. [46], Gimeno [47] they reported that decreased feed intake along with increased liver weights in birds is consistent with earlier reports. Bautista et al. [48] concluded that an average AF concentration of 0.95 mg/kg reduced both feed intake and daily weight gain by 11 percent, and worsened feed conversion by 6 percent. In the affected bird's metabolic changes lead to enlargement of liver, kidney and spleen as well as a decrease in the size of bursa of Fabricius, thymus and testes. With high dose exposure, fat accumulates inside the cell of the liver as clear vacuoles. As a result, the liver is greatly enlarged, yellow and friable (easily broken). There is general agreement that dietary AF reduces weight gain and feed intake, and worsens feed efficiency. Previous research indicated that the reduced growth rate because of AF ingestion in the diet is primarily due to the reduction in feed intake. Bautista et al. [48], Cheng et al. [49] found that, aflatoxicosis represents one of the serious diseases of poultry, livestock and other animals. *Aspergillus flavus* is known to produce a group of extremely toxic metabolites, of which aflatoxin B₁ (AFB₁) is most potent. Avian species especially chickens; goslings, ducklings and turkey poult are most susceptible to AFB toxicity. External examination of carcass did not show any disease changes. On post-mortem examination, the carcass

revealed large cirrhotic pale friable liver with perihepatitis. Heart, Lungs, Spleen and kidney were normal. Salpingitis, impaction of the oviduct with cheesy material and egg peritonitis were also noted. The case was diagnosed as aflatoxicosis. Case reports of aflatoxicosis in ducks are rare. Chronic exposure to low levels of aflatoxin can result in cancer and immunosuppression. Decreased feed intake along with increased liver weights in birds is consistent with earlier reports. Even though prevention and avoidance are the best way to control aflatoxicosis, natural contamination of crops with *A. flavus* is some times unavoidable. Andretta et al. [46], Wan et al. [19] Aflatoxin at a dietary concentration of 1 mg/kg or more caused severe reductions in growth and immune response in broilers, whereas 2 mg/kg increased the relative weight of the liver and decreased the relative weight of the bursa of Fabricius.

3.5 Improve Kidney Weights of Ducklings Given Feed Contaminated with aflatoxin B₁

Effect of aflatoxin B₁ exposure on average Kidney weight of duckling and its control were studied. Data were tabulated in Table 4 the presented data showed that enlargement in kidney weight in contaminated groups. Aflatoxin B₁ alone was found to significantly (P < 0.01 & P < 0.05) increase average kidney weight of all tested ducklings compared with the control group (Un-treated). It was increased from 4.5 g to 6.9 g when ducklings were given feed contaminated with 0.018 ng /100 g of AFB₁. Whereas, all ducklings received Propolis extracts plus aflatoxin B₁ were improved and appeared comparable to the control. Propolis extract was found significantly (P < 0.01 & P < 0.05) improve




Fig. 2. Showing enlarged kidneys of duckling treated with only aflatoxin (group 5) in relation to control one (G 6).

Fig. 3. Showing enlarged liver and conjunction of duckling treated with only aflatoxin (group 5) compared to control one (G 6)



Fig. 4. Showing small size gall bladder in control group and enlarged gall bladder of duckling treated with aflatoxin plus propolis group 2 (5% conc.) as well as group 3 (10%conc.) in relation to control one (Control G6)

all kidney weight. Continuous improve kidney weight with increasing the concentration of propolis extract at 2%, 5%, 10%. This improves was 6 g in the average kidney with 2%. While at 5% it gives 5.70 g , at 10% it gives 5.6 g Enhanced average kidney weight was 5.5 g which recorded with 20% propolis as compared with untreated group (Control G6).

3.6 Histological Studies

The biochemical results reported in the current study were confirmed by the histological results. Microscopic examination of liver and kidney tissues showing, changes as well as enhanced of histological tissues were confirmed and photographed as shown in Figs (1-38).

3.6.1 For control ducklings G 6 (free aflatoxin B₁)

No histopathological changes were detected in the examined livers and kidneys of these ducklings. The liver showed normal histological criteria of blood vessels, bile ducts and hepatic

cords. Rarely small numbers of hepatocytes showed vacuolar and hydropic degeneration (Fig. 5a). The kidneys revealed the normal histological appearance of both cortex and medulla. The renal cortex showed the normal histological structure of glomeruli and convoluted tubules (Fig. 5b).

Table 4. Improve average kidney weight of ducklings exposed aflatoxin B₁ by using propolis extract

Groups	Kw.	I	% I	%R
G1	6.00	1.5	0.25	99.75
G2	5.70	0.73	16.2	83.8
G3	5.60	1.1	19.6	80.4
G4	5.50	1	18.2	81.8
G5	6.90	2.4	34.8	65.2
G6 (control)	4.50			
L.S.D@1%	0.581			
L.S.D@5%	0.381			

*Kw.=kidney weight; I=increased; % I =increased percent; %R=Reduction




Fig. 5a. Liver of control duckling, and b showing normal histological criteria of hepatic tissue with vacuolar and hydropic degeneration of some hepatocytes (arrow). H&E stain x 200

Fig. 5b. kidney of control duckling, Group 6, showing normal histological criteria of glomeruli (G) and renal tubules (T). H & E stain x 200

3.6.2 Effect of aflatoxin B₁ alone on liver and kidney tissue of ducklings (G 5)

Ducklings received feed contaminated with 0.018 ng/ml aflatoxin B₁ at rate 1 ml/100 g feed resulted that, the livers showed congestion of the central veins and portal blood vessels. Fibrin thrombi were noticed in the lumen of the central and portal veins (Fig. 6). Activation of Von Kupffer,s cells and the presence of small numbers of macrophages and lymphocytes were noticed in the dilated sinusoids (Fig. 7). Perivascular oedema mixed with inflammatory cellular infiltration mainly lymphocytes (Fig. 8) was occasionally observed. Multifocally, the perivascular interstitium was expanded by mononuclear cellular aggregates (Fig.9) mainly macrophages and lymphocytes (Fig. 10). Similar areas of mononuclear cellular aggregation displaced the hepatocytes were seen. Diffusely, the hepatic parenchyma was distorted by marked

vacuolar and hydropic degeneration of the hepatocytes characterised by swollen, pale, vacuolated cytoplasm (Fig. 11). Focal areas of coagulative necrosis characterised by retention of hepatic cord architecture and shrunken hepatocytes with homogenous eosinophilic cytoplasm and pyknotic nuclei. Occasionally, lytic necrosis of the hepatocytes characterised by loss of hepatic cord architecture and replaced by erythrocytes, fibrin and small numbers of inflammatory cells were noticed (Fig.12). Moreover, the portal areas revealed mononuclear cellular infiltration and hyperplasia of the biliary epithelium.

The examined kidneys revealed congestion of the cortical blood vessels and intertubular capillaries (Fig. 13). There were aggregates of mononuclear inflammatory cells, mainly macrophages and lymphocytes, separating surrounding and effacing variable areas of renal




Fig. 6. Liver of duckling treated with aflatoxin, Group 5, showing marked dilatation and congestion of portal vein (PV) with fibrin thrombus (Th) in the lumen. H&E stain x 200.

Fig. 7. Liver of duckling treated with aflatoxin, Group 5, showing small numbers of macrophages (arrow) and lymphocytes (arrow head) in the dilated sinusoids. H&E stain x 400.

Fig. 8. Liver of duckling treated with aflatoxin, Group 5, showing perivascular oedema (asterisk) mixed with lymphocytic cellular infiltration (arrowhead). H&E stain x 200.




Fig. 9. Liver of duckling treated with aflatoxin, Group 5, showing perivascular mononuclear cellular aggregates (asterisk). H&E stain x 200

Fig. 10. High power of the previous figure showing perivascular mononuclear cellular aggregates mainly macrophages (arrow) and lymphocytes (arrow head). H&E stain x 400

Fig. 11. Liver of duckling treated with aflatoxin, Group 5, showing marked vacuolar and hydropic degeneration of the hepatocytes characterised by swollen, pale, vacuolated cytoplasm (arrowhead). H&E stain x 400

Fig. 12. Liver of duckling treated with aflatoxin, Group 5, showing lytic necrosis (N) of the hepatocytes characterised by loss of hepatic cord architecture and replaced by erythrocytes (arrow), fibrin and small numbers of inflammatory cells (arrowhead). H&E stain x 400




Fig. 13. Kidney of duckling treated with aflatoxin, Group 5, showing congestion of the cortical blood vessels (V) and intertubular capillaries (C). H&E stain x 100

Fig. 14. Kidney of duckling treated with aflatoxin, Group 5, showing aggregates of mononuclear inflammatory cells (M), separating, surrounding and effacing variable areas of renal architecture in the cortex. H&E stain x 400

Fig. 15. Kidney of duckling treated with aflatoxin, Group 5, showing enlarged glomeruli with degeneration of glomerular tufts characterised by vacuolisation (arrowhead) of mesangial/endothelial cells. H&E stain x 400

architecture in the cortex (Fig. 14). The glomeruli were enlarged and revealed either degeneration of glomerular tufts characterised by vacuolisation of mesangial/endothelial cells (Fig. 15) or coagulative necrosis with retention of tuft architecture, hypereosinophilic cytoplasm and pyknosis (Fig. 16). Multifocally, there was vacuolar and hydropic degeneration of the lining epithelial cells of some proximal and distal convoluted tubules characterised by swollen pale

vacuolated cytoplasm (Fig. 17). Coagulative necrosis of tubular epithelial cells characterised by hypereosinophilic cytoplasm and pyknotic nuclei were also detected (Fig. 18). The lumina of some renal convoluted tubules contained homogenous, brightly eosinophilic material (hyaline casts) (Fig. 19). Moreover, remaining tubules at all levels were moderate to markedly ectatic and lined by attenuated epithelium.




Fig. 16 Kidney of duckling treated with aflatoxin, Group 5, showing coagulative necrosis of glomerular tuft cells with retention of tuft architecture, hypereosinophilic cytoplasm and pyknosis(arrowhead). H&E stain x 400

Fig. 17 Kidney of duckling treated with aflatoxin, Group 5, showing marked vacuolar and hydropic degeneration of the lining epithelial cells of some proximal and distal convoluted tubules characterised by swollen pale vacuolated cytoplasm(arrowhead). H&E stain x 400

Fig. 18 Kidney of duckling treated with aflatoxin, Group 5, showing coagulative necrosis of tubular epithelial cells characterised by hypereosinophilic cytoplasm and pyknotic nuclei (arrowhead). H&E stain x 400

Fig. 19 Kidney of duckling treated with aflatoxin, Group 5, showing hyaline casts (Hc) in the lumen of some renal convoluted tubules. H&E stain x 400.

7. IMPROVE LIVER AND KIDNEY TISSUES BY USING PROPOLIS EXTRACT

7.1 Ducklings Received Aflatoxin B₁ Plus Propolis 2% (G 1)

The livers of the treated ducklings showed congestion of the central veins, portal blood vessels. Perivascular edemas infiltrated with mononuclear inflammatory cells were noticed in few cases. There were fibrin thrombi partially or completely occluded the lumen of the portal veins; the portal areas revealed a mild hyperplastic proliferation of the bile duct epithelium (Fig. 20). Multifocally, the surrounded hepatocytes exhibited mild to moderate degeneration characterised by discrete pale variable sizes cytoplasmic vacuoles (Fig. 21).

Occasionally, individual coagulative necrosis of the hepatocytes hypereosinophilic cytoplasm and pyknosis shrunken hepatic cells with hypereosinophilic cytoplasm and pyknotic nuclei were also noticed (Fig. 22).

The examined kidneys revealed congestion of the renal blood vessels and intertubular capillaries. The renal cortex revealed perivascular oedema mixed with few numbers of lymphocytes. Rarely, glomerular tufts exhibited necrosis of mesangial/endothelial cells characterised by hypereosinophilic cytoplasm and pyknosis Multifocally, the cortical renal tubules were lined by swollen, and vacuolated epithelial cells (degeneration) (Fig. 23) or occasionally by epithelial cells with hypereosinophilic cytoplasm and pyknotic nuclei (necrosis) (Fig. 24).




Fig. 20. Liver of duckling treated with aflatoxin and propolis at 2%, Group 1, showing fibrin thrombus (Th) in the lumen of the portal vein and the mild hyperplastic proliferation of the bile duct epithelium (arrowhead). Note also cytoplasmic vacuoles in the hepatocytes. H&E stain x 400

Fig. 21. Liver of duckling treated with aflatoxin plus propolis extract at 2%, Group 1, showing individual coagulative necrosis of the hepatocytes with hyper eosinophilic cytoplasm and pyknotic nuclei (arrowhead). H&E stain x 400

Fig. 22. Kidney of duckling treated with aflatoxin plus propolis extract at 2%, Group 1, showing vacuolar and hydropic degeneration (D) of renal tubules. H&E stain x 200




Fig. 23 High power of the previous figure showing degenerated renal tubules lined by swollen, vacuolated epithelial cells (arrowhead). H&E stain x 400

Fig. 24 Kidney of duckling treated with aflatoxin plus propolis extract at 2%, Group 1, showing necrotic renal tubules lined by epithelial cells with hyper eosinophilic cytoplasm and pyknotic nuclei (arrowhead). H&E stain x 400

7.2 Ducklings Received Aflatoxin B₁ plus Propolis 5% (G 2)

The livers showed congestion of the portal blood vessels and leucocytic cellular infiltration in the portal areas particularly lymphocytes (Fig. 25). Fibrin thrombi partially or completely occluded the lumen of the portal veins were attached to the blood vessel wall (Fig. 26). Multifocally, degenerative changes of the hepatocytes in the form of vacuolar and hydropic degeneration were commonly observed around central veins (Fig. 27). Occasionally, individual to random areas of coagulative necrosis characterised by retention of hepatic cord architecture and shrunken hepatocytes with hyper eosinophilic cytoplasm

and pyknotic nuclei were noticed (Fig. 28). The portal areas revealed mild hyperplasia of the biliary epithelium. Furthermore, periductal infiltrate by moderate numbers of lymphocytes was also detected (Fig. 29).

The examined kidneys revealed congestion of the cortical blood vessels and intertubular capillaries with focal areas of intertubular haemorrhages (Fig. 30). Rarely, the blood vessels revealed hypertrophy and vacuolar degeneration of the wall. Multifocally, the proximal and distal convoluted tubules in the renal cortex showed either coagulative necrosis or degenerative changes characterised by cloudy swelling, vacuolar and hydropic degeneration of

their lining epithelium (Fig. 31). Necrotic tubular epithelial cells were hyper eosinophilic with shrunken, pyknotic nuclei, loss of cellular detail, and karyolysis (Fig. 32). Degenerated tubular epithelial cells were occasionally swollen with pale vacuolated cytoplasm (Fig.33).

7.3 Ducklings Received Aflatoxin B₁ Plus Propolis 10% (G 3)

The livers of the treated ducklings showed congestion of the central veins and portal blood

vessels, with mild perivascular inflammatory cellular infiltration mainly lymphocytes (Fig. 34). Multifocally, there were random areas of vacuolar and hydropic degeneration of the hepatocytes characterised by swollen pale vacuolated cytoplasm (Fig. 35). Occasionally, individual coagulative necrosis of the hepatocytes characterised by retention of hepatic cord architecture and shrunken hepatocytes with hyper eosinophilic cytoplasm and pyknotic nuclei were noticed.




Fig. 25. Liver of duckling treated with aflatoxin and propolis 5%, Group 2, showing individual coagulative necrosis of the hepatocytes with hyper eosinophilic cytoplasm and pyknotic nuclei (arrow). H&E stain x 400




Fig. 26. Liver of duckling treated with aflatoxin plus propolis extract at 5%, Group 2, showing congestion of the portal blood vessels (V) and lymphocytic cellular infiltration (L) in the portal areas. H&E stain x 400




Fig. 27. Liver of duckling treated with aflatoxin plus propolis extract at 5%, Group 2, showing fibrin thrombus (Th) were attached to the wall of the portal vein (Pv). H&E stain x 100




Fig. 28. Liver of duckling treated with aflatoxin plus propolis extract at 5%, Group 1, showing vacuolar and hydropic degeneration (arrow) of the hepatocytes around central veins (CV). H&E stain x 200




Fig. 29. Liver of duckling treated with aflatoxin extract plus propolis at 5%, Group 2, showing individual coagulative necrosis of the hepatocytes with hyper eosinophilic cytoplasm and pyknotic nuclei (arrow). H&E stain x 400




Fig. 30. Kidney of duckling treated with aflatoxin plus propolis extract at 5%, Group 2, showing congestion of the cortical blood vessels (V) and intertubular capillaries with focal areas of intertubular haemorrhages (H). H&E stain x 100




Fig. 31. Kidney of duckling treated with aflatoxin plus propolis extract at 5%, Group 2, showing coagulative necrosis (N) and degenerative changes (D) of the lining epithelium of proximal and distal convoluted tubules in the renal cortex. H&E stain x 200

Fig. 32. Kidney of duckling treated with aflatoxin plus propolis extract at 5%, Group 2, showing necrotic tubular epithelial cells with shrunken, pyknotic nuclei (arrowhead), loss of cellular detail, and karyolysis (arrow). H&E stain x 400

Fig. 33. Kidney of duckling treated with aflatoxin plus propolis extract at 5%, Group 2, showing degenerated tubular epithelial cells with swollen pale vacuolated cytoplasm (arrowhead). H&E stain x 400. cytoplasm

The examined kidneys revealed congestion of the renal blood vessels and intertubular capillaries with focal areas of intertubular haemorrhages (Fig. 36). The glomeruli and renal convoluted tubules in both cortex and medulla revealed normal histological appearance similar

to the control group. It was an enhancement as shown in (Fig. 37). Rarely, few tubules were lined by swollen, vacuolated epithelial cells (degeneration) or by epithelial cells with hypereosinophilic cytoplasm and pyknotic nuclei (necrosis). It was the best concentration used.




Fig. 34. Liver of duckling treated with aflatoxin plus propolis extract at 10%, Group 3, showing congestion of the portal blood vessels (Pv), with mild perivascular inflammatory cellular infiltration, mainly lymphocytes (L). H&E stain x 400

Fig. 35. Liver of duckling treated with aflatoxin plus propolis extract at 10%, Group 3, showing random areas of vacuolar and hydropic degeneration (D) of the hepatocytes characterised by swollen pale vacuolated cytoplasm. H&E stain x 200

Fig. 36. Kidney of duckling treated with aflatoxin plus propolis extract at 10%, Group 3, showing congestion of the renal blood vessels (V) and intertubular capillaries with focal areas of intertubular haemorrhages (H). H&E stain x 200

Fig. 37. Kidney of duckling treated with aflatoxin plus propolis extract at 10%, Group 3, showing the normal histological appearance of the glomeruli (G) and renal convoluted tubules (T) in the cortex. H&E stain x 200




Fig. 38. Liver of duckling treated with aflatoxin plus propolis extract at 20%, Group 4, showing mild vacuolar and hydropic degeneration of the hepatocytes characterised by swollen pale vacuolated cytoplasm (arrow head). H&E stain x 200

Fig. 39. Liver of duckling treated with aflatoxin plus propolis extract at 20%, Group 4, showing massive areas of haemorrhages (H) around the congested blood vessel (V). H&E stain x 100

Fig. 40. Liver of duckling treated with aflatoxin plus propolis extract at 20%, Group 4, showing massive areas of haemorrhages (H) in between the convoluted tubules. H&E stain x 200

Fig. 41. Kidney of duckling treated with aflatoxin plus propolis extract at 20%, Group 4, showing the normal histological appearance of renal convoluted tubules (T) in the cortex. H&E stain x 400

7.4 Ducklings Received Aflatoxin B₁ Plus Propolis 20% (G 4)

The microscopical examination of livers showed congestion of the central veins and portal blood vessels. Multifocally, mild vacuolar and hydropic degeneration of the hepatocytes characterised by swollen pale vacuolated cytoplasm were seen (Fig. 38). However, variable areas of hepatic parenchyma preserve normal histological criteria as a result of liver improvement.

The examined kidneys revealed congestion of the renal blood vessels and intertubular capillaries. The renal cortex revealed massive areas of haemorrhages around the congested blood vessel (Fig. 39) and in between the convoluted tubules (Fig. 40). Moreover, the glomeruli and renal convoluted tubules in both cortex and medulla revealed normal histological appearance similar to the control group (Fig. 41). It was improved exhibit the renal cortex revealed massive areas of haemorrhages around the congested blood vessel.

Similar results were obtained by many investigators. Bibu [28], Gimeno [47] reported that one to seven day old duckling given a feed with a concentration of 300 to 600 ppb of AFB₁ for 7 to 14 days had serious hepatic lesions, as well as significant death rate. Bautista et al. [48] concluded that, the toxicity of aflatoxins, in general, may be categorised as acute or chronic. Acute aflatoxicosis causes marked signs of

disease or death. The liver is usually pale, atrophied or necrotic. Symptoms include loss of appetite and lethargy leading to death. Chronic aflatoxicosis are not readily discernible in affected animals. Visible symptoms may include reduced growth, appetite, and feed efficiency. The liver may appear normal but histological examination will likely reveal abnormalities. Aflatoxins had detrimental effects, which increased with level, on phagocytotic ability and tumour necrosis factor-like substance secretion [49] and caused liver damage [50] in ducks. AF acts as an inhibitor of protein synthesis and, subsequently, dividing cells and tissues with a high protein turnover such as that found in the liver, immune system or gut epithelium, which is most susceptible to the toxic effects of AF. In this respect, exposure to AF has been demonstrated to suppress the immune response in poultry. AF can repress the development of the thymus gland or influence the relative weight of the bursa of Fabricius, which may result in serious deficiencies in both cellular and antibody responsiveness of the chicken immune system [51].

8. CONCLUSION

This study is one of the first where an antimicrobial property of propolis against *A. flavus* fungus which contaminant chicken and fish feeds. These results suggest that the propolis extract is excellent antifungal activities. This approach is considered as an environmental

friendly approach in contrast to physical and biological techniques of detoxification. Various methods of decontamination are found to vary in their efficacy in toxin removal from feed products. The cost of the decontamination process is very important in choosing the cheapest and the most effective method in aflatoxin removal from the contaminated feed products.

CONSENT

It is not applicable.

ETHICAL APPROVAL

As per international standard or university standard ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Pittet A. Natural occurrence of mycotoxins in foods and feeds – an updated review, *Rev. Méd. Vét.* 1998;149:479–492.
2. Vasanthi S, Bhat RV. Mycotoxins in foods-occurrence, health and economic significance and food control measures. *Ind. J. Med. Res.* 1998;108:212-224.
3. Fink-Gremmels J. Mycotoxins in cattle feeds and carryover to dairy milk: A review. *Food Addit. Contam.* 2008;25(2):172-180.
4. Wu J, Miller D, Cayman EA. Bt maize and mycotoxin reduction: Economic impacts in the United States and the developing world. *J. Toxicol. Toxin Rev.* 2004;23:397-424.
5. Upadhaya SD, Park MA, Jong KHA. Review: Mycotoxins and their biotransformation in the rumen. *Asian-Aust. J. Anim. Sci.* 2010;23(9):1250–60.
6. Krnjaja V, Stojanović Lj, Cmijanić R, Trenkovski S, Tomašević D. The presence of potentially toxigenic fungi in poultry feed. *Biotechnology in Animal Husbandry.* 2008;24(5-6):87-93.
7. Anonymous. Feed quality problems and management strategies. Available:<http://repository.seafdec.org.ph> on July 31, 2017 at 8:04 PM CST; cited from Cruz PS. Fish nutrition and feeds '94 proceedings. Proceedings of the National Seminar-Workshop on Fish Nutrition and Feeds; SEAFDEC Aquaculture Department, Iloilo, Philippines. 1996;64-73.
8. Dalvi RR. An overview of aflatoxicosis of Poultry: Its characteristics, prevention and reduction. *Vet. Res. Commun.* 1986;10(1): 429-443.
9. Silici S, Unlu M, Vardar-Unlu G. Antibacterial activity and phytochemical evidence for the plant origin of Turkish propolis from different regions. *World J Micro Biot.* 2007;23:1797-1803.
10. Simone-Finstrom M, Spivak M. Propolis and bee health: The natural history and significance of resin use by honey bees. *Apidologie.* 2010;41:295-311.
11. Mavri A, Abramovic H, Polak T, Bertocelj J, Jamnik P, Mozina SS, Jersek B. Chemical properties and antioxidant and antimicrobial activities of Slovenian propolis. *Chem Biodivers.* 2012;9:1545-1558.
12. Ammar MI, El-Naggar MA. Screening and characterization of fungi and their associated Mycotoxins in some Fruit Crops. *International Journal of Advanced Research.* 2014;2(4):1216-27.
13. Özdemir AE, Çandır EE, Kaplankiran M, Soyulu EMN, Şahinler A. Gül. The effects of ethanol-dissolved propolis on the storage of grapefruit cv. Star Ruby. *Turkish Journal of Agriculture and Forestry.* 2010;34:155-162.
14. Ngoepe EC, Straker C. Propolis as a natural antimicrobial agent for control of fungal pathogens of plants. Ph. D. Honours dissertation submitted to the University of the Witwatersrand, South Africa; 2004.
15. Giovanelli LC. Evaluation of an ethanolic extract of propolis as a potential pre- and post-harvest fungicide for 'Fuerte' avocado (*Persea americana* Mill.) fruits and orchards. M.Sc. degree, Faculty of Science, University of the Witwatersrand, Johannesburg. 2008;113.
16. Embaby EM, Nahed M. Ayaat, Mona M. Abd El-Galil, Nassr-Allah Abdel-Hamid, Mona M. Gouda. Mycoflora and mycotoxin contaminated chicken and fish feeds. *Middle East Journal of Applied Sciences.* 2015;05(04):1044-54.
17. Mehrim AI, Salem MF. Medicinal herbs against aflatoxicosis in Nile tilapia (*Oreochromis niloticus*): Clinical signs, postmortem lesions and liver histopathological changes. *Egy. J. aquac.* 2013;3(1): 13-25.
ISSN: 2090-7877

18. NRC. Nutrient requirements of poultry. 9th rev. ed. Natl. Acad. Press, Washington, DC; 1994.
19. Wan XL, Yang ZB, Yang WR, Jiang SZ, Zhang GG, Johnston SL, Chi F. Toxicity of increasing aflatoxin B₁ concentrations from contaminated corn with or without clay adsorbent supplementation in ducklings. *Poultry Science*. 2013;92:1244–53.
20. Barham D, Trinder P. An improved colour reagent for the determination of blood glucose by the oxidase system. *Analyst*. 1972;97(151):142-45.
21. Bartels H, Bohmer M, Heirli C. Serum creatinine determination without protein precipitation. *Clin. Chem. Acta*. 1972;37: 193–197.
22. Drury AA, Wallington EA. Carleton's histological technique. 5th Ed., Oxford University press, New York, Toronto; 1980.
23. Saddiq AA, Kalifa SA. Impact of fungal content of some Arabic nuts to induce kidney toxicity and agonistic action of natural resources. *African Journal of Microbiology Research*. 2011;5(9):1046-1056.
24. Omar N. Effect of some aflatoxins on a lymphatic organ (spleen) of male albino rats (Histopathological Study) .*The Egyptian Journal of Hospital Medicine*. 2002;48:357–367.
25. Snedecor GW, Cochran WG. Statistical methods. 7th Ed. Iowa state Univ. Press. Ames Iowa, USA; 1989.
26. Duncan DB. Multiple rang and multiple F test. *Biometrics*. 1955;11:1-42.
27. Gimeno A. Los hongos y las Micotoxinas en la Alimentacion Animal ;Conceptos, Problemas, control y Recomendaciones. 2000;1-49.
Available:www.engormix.com(lra:mictotoxinas.Seccionenespanol)(consultado
28. Bibu KJ. Aflatoxicosis in a white pekin duck. *Veterinary World*. 2011;4(5):215.
Available:www.veterinary world .org 25-03
29. Mariana VG, María LF, Silvia LRG, Pardo AG, Pose GN. Mycotoxins and mycotoxigenic fungi in poultry feed for food-producing animals. *The Scientific World Journal*. 2014;9. [Article ID 968215]
30. Gauthier J, Ludlow R. Chicken health for dummies, poultry feeds and feeding. *Poultry Feeds and Feeding. Hand Book, Canadian Poultry Consultants Ltd*. 2016; (Chapter II):1-877-449-3447.
31. Wu J, Miller D, Casman EA. Bt maize and mycotoxin reduction: Economic impacts in the United States and the developing world. *J. Toxicol. Toxin Rev*. 2004;23:397-424.
32. Yunus AW, Razzazi-Fazeli E, Bohm J. Aflatoxin B1 in affecting broiler's performance, immunity, and gastrointestinal tract: A review of history and contemporary issues. *Toxins*. 2011;3:566-590.
33. Bintvihok A. Controlling aflatoxin danger to duck and duck meat. *World Poultry Sci J*. 2001;17:18-20.
34. Anjum AD. Outbreak of infectious bursal disease in vaccinated chickens due to aflatoxicosis. *Indian Journal of Animal Science*. 1994;71:322-324.
35. Verma D. New facts about mycotoxin control-intensive research in the field of mycotoxin deactivation gives new insights! *BioMin India*; 2007.
36. Chen X, Grenier B, Applegate TJ. Applegate aflatoxins in poultry. *Purdue Animal Sciences Purdue extension*. 2013; 6.
Available:www.ag.purdue.edu/ANSC, www.Extension.Purdue.Edu
37. Mabbett T. Keep feeds free from fungi. *African Farming*. View at Google Scholar. 2004;15–16.
38. Radziejewska RC, Stuper K, Szablewski T. Microflora and mycotoxin contamination in poultry feed mixtures from western Poland. *Annals of Agricultural and Environmental Medicine*. 2013;20(1):30–35.
39. Greco (Mariana) V, Franchi (María) L, Golba (Silvia) LR, Pardo AG, Pose GN. Mycotoxins and mycotoxigenic fungi in poultry feed for food-producing animals. *The Scientific World Journal*. 2014;9. [Article ID 968215]
40. Chang, W.; Xie, Q.; Zheng, A.; Zhang, S.; Chen, Z.; Wang, J.; Liu, G. and Cai, H. Effects of aflatoxins on growth performance and skeletal muscle of Cherry Valley meat male ducks. *Animal Nutrition*. 2016; 2, 186-191.
41. Akande KE, Abubakar MM, Adegbola TA, Bogoro SE. Nutritional and health implications of mycotoxins in animal feeds: A review. *Pakistan J. Nutr*. 2006;5(5):398-403.
42. Sultana N, Hanif NQ. Mycotoxin contamination in cattle feed and feed ingredients. *Pakistan Vet. J*. 2009;29(4): 211-213.

43. Grenier B, Applegate TJ. Modulation of intestinal functions through the mycotoxin ingestion. *Toxins*; 2012.
44. Ortatatli M, Oguz H. Ameliorative effects of dietary clinoptilolite on pathological changes in broiler chicken during aflatoxicosis. *Res .Vet. Sci.* 2002;71:59-66.
45. Verma J, Johri TS, Swain BK, Ameena S. Effect of graded levels of aflatoxin, ochratoxin and their combinations on the performance and immune response of broilers. *British Poultry Science.* 2004;45: 512-518.
46. Andretta I, Kipper M, Lehnen CR, Hauschild L, Vale MM, Lovatto PA. Meta-analytical study of productive and nutritional interactions of mycotoxins in broilers. *Poultry Science.* 2011;90:1934-1940.
47. Gimeno A. Revision generica del problema de los hongos y las micotoxinas en la alimentacion animal. 1999;1-53. Available:www.mycotoxin.com
48. Bautista MN, Lavilla-Pitogo CR, Subosa PF, Bagino ET. Aflatoxin B1 contamination of shrimp feeds and its effect on growth and hepatopancreas of pre-adult *Penaeus monodon*. *J. Sci. Food Agric.* 1994;65:5-11.
49. Cheng YH, Shen TF, Chen BJ. Induction of changes in morphology, reactive nitrogen/ oxygen intermediates and apoptosis of duck macrophages by aflatoxin B1. *Asian-australas. J. Anim. Sci.* 2002;11:1639–1645.
50. Ostrowski-Meissner HT. Effect of contamination of diets with aflatoxins on growing ducks and chickens. *Trop. Anim. Health Prod.* 1983;15:161–168.
51. Celik I, Oguz H, Demet O, Donmez HH, Boydak M, Sur E. Efficacy of polyvinylpyrrolidone in reducing the immunotoxicity of aflatoxin in growing broilers. *British Poultry Science.* 2000;41:430-439.

© 2018 Ayaat et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/27920>*